

Large Games with Heterogeneous Players

Ricardo Serrano-Padial*

August 18, 2020

Abstract

We study large games played by heterogeneous agents whose payoffs depend on the aggregate action and provide novel equilibrium selection and comparative statics results. We prove the equivalence between the global games selection and potential maximization in games with strategic complementarities, and characterize the selected equilibrium as the solution to maximizing the ex-ante expected payoffs of a player with pessimistic beliefs. To obtain our results we uncover two key properties. First, we show that potential games exhibit quasilinear payoffs, which include as examples Cournot competition, public goods and externalities, search models and coordination games. Second, we show that beliefs in the global game satisfy a generalized Laplacian property, which links average beliefs about the aggregate action to the uniform distribution. Our comparative statics results rely on average rather than pointwise conditions on payoffs and can be used to analyze both parameter changes and the impact of heterogeneity.

Keywords: aggregative games, potential games, global games, comparative statics, heterogeneity, noise-independent selection, Laplacian beliefs

1 Introduction

Large games in which individual payoffs depend on the aggregate behavior in the population are ubiquitous in economics and finance.¹ An incomplete list includes

*School of Economics, Drexel University; rspadial@gmail.com. I am deeply grateful to Ryota Iijima, Stephen Morris, Guillermo Ordoñez, Daisuke Oyama, Joel Sobel and Dai Zusai for helpful comments, as well as audiences at Academia Sinica, Hitotsubashi, Kyoto, UPenn and PETCO 2018 at Penn State.

¹These games are referred in the literature as (linear) aggregative games (Jensen, 2018), average-action games (Morris and Shin, 2003), games with aggregation (Dubey et al., 2006) or aggregate games (Martimort and Stole, 2012).

models of market competition (aggregate output), macroeconomic coordination (average search effort), public goods and externalities (sum of contributions), technology adoption (average investment), as well as binary-action games where payoffs depend on the fraction of agents adopting each action such as platform choice, bank runs, currency crises or games of regime change.

In all these economic phenomena, agent heterogeneity is a defining characteristic of the environment (e.g., agents may differ in costs, preferences, productivity or endowments), and understanding its effects is an integral part of the analysis. However, the introduction of heterogeneity complicates the three alternative ways in which equilibrium analysis is conducted, which involve imposing conditions that guarantee equilibrium uniqueness, selecting a particular equilibrium in the presence of multiplicity, or applying comparative statics results to the set of equilibria. For instance, heterogeneity limits the appeal of popular selection rules, such as those based on introducing incomplete information (e.g., global games), due to the inability to characterize the selected equilibrium or because of the lack of clear economic content behind the selection. In addition, most of the existing robust comparative statics results rely on monotonicity restrictions at the individual level, which may not apply to models where agents have divergent interests.

This paper addresses these limitations by providing new equilibrium characterization and comparative statics results for large games with heterogeneous payoff types. Specifically, we establish the equivalence between two commonly used equilibrium selection rules in large games with strategic complementarities: the global games selection (Carlsson and van Damme, 1993; Frankel et al., 2003) and potential maximization (Monderer and Shapley, 1996). Importantly, we characterize the selected equilibrium and give economic content to the selection by showing that maximizing potential coincides with maximizing the ex-ante payoffs of an agent with marginal beliefs who thinks that she is pivotal type, i.e., the last type contributing to the aggregate action. In addition, we generalize existing results on monotone comparative statics based on conditions that all types must satisfy by proposing weaker restrictions that only apply on average. In particular, we identify conditions on average payoffs under which the set of equilibrium aggregate actions ‘moves up’ after a change in parameters or in the distribution of types.

The games we study have payoffs that depend on the player’s action $a \in \mathbb{R}$, the aggregate action \bar{a} and the player’s type w . We obtain the equilibrium charac-

terization result by uncovering two key properties. First, we determine the payoff structure of potential games. Second, we pin down average beliefs in global games with heterogeneous payoffs.

Section 3 shows that potential games exhibit *quasilinear payoffs*, which take on the following form: $U(a, \bar{a}, w) = au(\bar{a}) + v(a, w)$. This payoff structure figures prominently in the economics literature, such as in the classic models of Cournot competition, Diamond’s search model, Tullock contests and most binary-action games, just to name a few.² Potential games are defined by the existence of a single function of the strategy profile (i.e., the potential function) that captures the change in individual payoffs of any player following a change in her strategy. Because the potential function must reflect individual payoff changes for all types, its existence imposes strong symmetry restrictions on the payoff impact of the aggregate action, which are only satisfied by quasilinear payoffs.

Section 4 presents a characterization of beliefs in global games, which we call the *Generalized Laplacian Property* since it generalizes to many-action, heterogeneous-player games both the Laplacian property of homogeneous binary-action games (Morris and Shin, 2003) and its counterpart for binary-action heterogeneous games (Sakovics and Steiner, 2012). In a global game, agents receive noisy signals about some payoff parameter and, because of this, face uncertainty about the aggregate action. The property states that the weighted average belief about the aggregate action is given by the uniform distribution, where the weights are proportional to the contribution of each type to the aggregate action.

Equipped with these properties, we show in Section 4 that changes in expected payoffs in the global game coincide with changes in potential as the noise vanishes, implying that the equilibrium in the global game converges to the strategy profile that maximizes potential. We characterize the selected equilibrium by pinning down the functional form of the potential function, and provide economic content behind the selection by deriving a dual representation of potential as the ex-ante payoffs

²Models with quasilinear payoffs include externalities (Dybvig and Spatt, 1983), technology adoption (Farrell and Saloner, 1985; Katz and Shapiro, 1986), contests (Tullock, 1980; Cornes and Hartley, 2005), common resources (Dasgupta and Heal, 1979), macroeconomic search (Diamond, 1982), cost-sharing (Moulin, 1990), bank runs (Diamond and Dybvig, 1983; Goldstein and Pauzner, 2005), currency crises (Obstfeld, 1986; Morris and Shin, 1998; Guimaraes and Morris, 2007), tax evasion (Bassetto and Phelan, 2008), regime change and coordination games (Sakovics and Steiner, 2012), crime waves (Bond and Hagerty, 2010), and blockchain (Abadi and Brunnermeier, 2018).

of a player with marginal beliefs. The proposed interpretation allows for a comparison with other selection rules such as Pareto dominance, which maximizes the ex-ante expected payoff given correct beliefs, and answers the open question about the meaning of maximizing potential in the context of aggregative games.³

In addition to our characterization result, we also present robust comparative statics for games with quasilinear payoffs that apply to the set of equilibrium aggregate actions (Section 5). We show that the set moves up following a change in parameters or in the distribution of payoff types if *average payoffs* across types satisfy a monotonicity restriction (increasing differences). In contrast, the existing literature relies on monotonicity restrictions on payoffs that must apply pointwise for all types (Milgrom and Shannon, 1994) or, alternatively, on monotonicity restrictions on best responses instead of on primitives of the game (Acemoglu and Jensen, 2010; Camacho et al., 2018). We obtain our results by recasting the problem of finding equilibrium as a fixed point problem over the set of aggregate actions instead of full strategy profiles. We highlight the differences with existing results by showing that the smallest and largest equilibrium average actions go up with parameters even if individual incentives are not monotone, that is, even if the game is not supermodular, which allows some agents to exhibit decreasing best responses. Accordingly, our results can be suitable for games where agents have diverging interests. Moreover, our comparative statics results regarding the distribution of types only rely on average restrictions on the heterogeneous component of payoffs $v(a, w)$, leading to a tractable analysis of the impact of heterogeneity on aggregate behavior. Independently of these results, our characterization of Nash equilibrium in terms of aggregate behavior could be useful in deriving conditions for equilibrium uniqueness.

Overall, the paper shows that in games with quasilinear payoffs one can obtain results by replacing pointwise conditions by average conditions on payoffs and beliefs. The paper contributions, which also include a novel definition of potential for games with continuous actions and types, touch upon many areas of economic theory. Accordingly, after presenting the main definitions and results, we discuss the related literature in Section 6.

³In their paper introducing potential games, Monderer and Shapley (1996) openly ask about the meaning of potential maximization (bottom of page 125, square brackets added for clarification):

“This raises the natural question about the economic content (or interpretation) of P^ [potential maximizer]: What do the firms [players] jointly try to maximize? We do not have an answer to this question.”*

2 Large Games

There is a continuum of players of measure one. Each agent is endowed with a type $w \in [\underline{w}, \bar{w}]$ and the mass of agent types is distributed in the population according to cdf F with density f . They simultaneously choose an action from the set $A \subset \mathbb{R}_+$, which can be any compact, countable union of single points and closed intervals. Leading examples include finite action and continuous action games. To simplify notation, we normalize actions so that the lowest action in A is set to zero.⁴ Let $a_{max} := \max A$ denote the highest action in A .

The payoffs of a player of type w choosing action a are given by $U(a, \bar{a}, \theta, w)$, where $\bar{a} \in [0, a_{max}]$ denotes the average of players' actions in the population and θ is a common parameter that belongs to the closed interval $\Theta \subset \mathbb{R}$. We assume that U is Lipschitz continuous and bounded in all its arguments.⁵ In addition, let

$$\Delta U(a, a', \bar{a}, \theta, w) = U(a, \bar{a}, \theta, w) - U(a', \bar{a}, \theta, w)$$

denote the payoff differences between choosing a and a' for a player of type w .

A strategy profile is given by a measurable mapping $\alpha : [\underline{w}, \bar{w}] \rightarrow A$. Let \mathcal{A} denote the set of measurable functions α .⁶ The average or aggregate action of players with types in any measurable subset $W \subseteq [\underline{w}, \bar{w}]$ under profile α is given by

$$\bar{\alpha}(\alpha, W) = \int_{\underline{w}}^{\bar{w}} \alpha(w) dF(w|w \in W), \quad (1)$$

Abusing notation we use $\bar{\alpha}(\alpha)$ to denote the aggregate action in the whole population $\bar{\alpha}(\alpha, [\underline{w}, \bar{w}])$. Accordingly, for any strategy profile $\alpha \in \mathcal{A}$, the payoffs of a player of type w are given by $U(\alpha(w), \bar{\alpha}(\alpha), \theta, w)$.

Formally, a game is given by the tuple $\Gamma = \{F, A, \theta, U\}$. Both F and θ are

⁴If $\min A \neq 0$ we can always redefine the set of actions to be $A' = \{a - \min A, a \in A\}$.

⁵By continuity in own action a we mean at interior points of A . Our results can be extended to payoffs that exhibit discontinuities at certain values of \bar{a} or θ , as is the case in regime change models (see [Serrano-Padial \(2018\)](#) for details).

⁶The restriction to measurable strategies ensures that payoffs are well defined. It is without loss of generality in supermodular games, which exhibit equilibrium strategies that are monotone functions from a measurable subset of \mathbb{R} to A and thus are measurable. However, it may be restrictive in general since it imposes that all agents of the same type use the same (pure) strategy.

common knowledge, i.e., Γ is a game of complete information. A Nash equilibrium (NE) of the game is a strategy profile α^* satisfying

$$\alpha^*(w) \in \arg \max_{a \in A} U(a, \bar{\alpha}(\alpha^*), \theta, w) \text{ for all } w \in [\underline{w}, \bar{w}].$$

3 Quasilinear Payoffs and Potential

In this section we show the equivalence between quasilinear payoffs and potential games. First, we define quasilinearity and potential.

Definition 1. Payoffs are *quasilinear* if there exist measurable functions u, v such that payoffs are given by

$$U(a, \bar{a}, \theta, w) = au(\bar{a}, \theta) + v(a, \theta, w),$$

or, more generally, if U can be expressed as $c(\theta, w)(au(\bar{a}, \theta) + v(a, \theta, w)) + u_0(\bar{a}, \theta, w)$ with $c(\theta, w) > \xi > 0$ for all θ, w .

Quasilinearity in binary-action games ($A = \{0, 1\}$) translates into payoff differences being *separable* in aggregate action and type, as defined in [Serrano-Padial \(2018\)](#). That is, $\Delta U(1, 0, \bar{a}, \theta, w) = u(\bar{a}, \theta) + v_\Delta(\theta, w)$, where $v_\Delta(\theta, w) = v(1, \theta, w) - v(0, \theta, w)$.

Most applications focus on continuous actions ($A = [0, a_{max}]$) or on binary actions (see [footnote 2](#)). Perhaps the leading example of continuous-action games with quasilinear payoffs and heterogeneous agents is the model of Cournot competition with heterogeneous cost functions, where a is individual output, u represents inverse demand and v are production costs. Binary action games with separable payoffs include most coordination games such as regime change models in which the probability of the regime failing is a function of the fraction attacking the regime.

We will use the following two examples featuring continuous actions to provide intuition about our results. The first one is about strategic complementarities in investment, and it will be used to illustrate how to derive the mentioned equilibrium selection rules and characterize the selected equilibrium. The second example is about negative externalities, e.g., due to congestion, and will highlight the usefulness of our comparative statics results.

Example 1 (Investment Game). Consider an economy populated by a continuum of heterogeneous firms choosing how much to invest in a new technology. Each firm chooses investment level $a \in [0, a_{max}]$. Unit returns on investment, given by $u(\bar{a}, \theta)$, are increasing in the degree of technology adoption \bar{a} and in the quality of the technology $\theta > 0$. Investment costs are quadratic and inversely proportional to the firm's productivity type $w \in [\underline{w}, \bar{w}] \subset \mathbb{R}_{++}$, which is distributed according to F in the population with $Ew = 1$. Specifically, payoffs are given by

$$U(a, \bar{a}, \theta, w) = au(\bar{a}, \theta) - \frac{a^2}{2} \frac{1}{w}. \quad (2)$$

Example 2 (Negative externalities). A continuum of agents choose their individual consumption level $a \in [0, a_{max}]$ of an exhaustible/congestible good (e.g., roads, cell-phone bandwidth, natural resources). Payoffs are given by the benefit from usage, which depends on type w and a common attribute θ of the good, minus (linear) costs, which increase with average consumption \bar{a} . Specifically,

$$U(a, \bar{a}, \theta, w) = b(a, \theta, w) - c(\bar{a}, \theta)a,$$

where b, c are differentiable, b is strictly concave in a and c is increasing in \bar{a} .

The definition of potential games was initially introduced for finite games, and involves the existence of a mapping from strategy profiles to the real line, the potential function, whose change after a single player switches strategies coincides with the player's payoff change. Since the game $\Gamma = \{F, A, \theta, U\}$ exhibits a continuum of players and types, we account for the fact that any single type switching strategies has no measurable impact on the strategy profile by defining potential as a function whose infinitesimal change after a type switches strategies coincides with the type's change in payoffs.⁷ We do so by using a mixture distribution that places a positive mass on agents of type w and take the limit of the change in potential as this mass goes to zero. Let $\delta(w)$ denote the Dirac delta distribution that places all the probability mass on type w and \mathcal{F} the space of distribution functions on $[\underline{w}, \bar{w}]$.

⁷Existing definitions of potential focus on finite games (e.g., [Monderer and Shapley, 1996](#)) or population games with a discrete distribution of types (e.g., [Sandholm, 2009](#)). We can accommodate finite types $\{1, \dots, N\}$ by setting $[\underline{w}, \bar{w}] = [0, 1]$ and $F = U[0, 1]$. Accordingly, we can partition $[0, 1]$ into N subintervals such that each interval i has length equal to the mass of discrete type i and assign the payoff function of discrete type i to all w in interval i .

Definition 2 (Potential). Let F_w^ε denote the mixture distribution $(1 - \varepsilon)F + \varepsilon\delta(w)$ for $\varepsilon \in (0, 1)$. The game Γ is a *potential game* if there exists a continuous functional $V : \mathcal{A} \times \mathcal{F} \times \Theta \rightarrow \mathbb{R}$ such that, for all w , all $a \in A$, and all $\alpha \in \mathcal{A}$

$$\lim_{\varepsilon \rightarrow 0} \frac{V(\alpha', F_w^\varepsilon, \theta) - V(\alpha, F_w^\varepsilon, \theta)}{\varepsilon} = \Delta U(a, \alpha(w), \bar{\alpha}(\alpha), \theta, w), \quad (3)$$

where $\alpha'(w) = a$ and $\alpha'(w') = \alpha(w')$ for all $w' \neq w$. The functional V is called the potential of the game.⁸

Γ is a *weighted potential game* if there exists a function $\psi(w, \theta) > \zeta > 0$ such that the game $\{F, A, \theta, \tilde{U}\}$ with payoffs given by $\tilde{U}(a, \alpha, \theta, w) = \psi(w, \theta)U(a, \alpha, \theta, w)$ is a potential game. The potential of $\{F, A, \theta, \tilde{U}\}$ is the weighted potential of Γ .

Our first result shows that quasilinearity is a necessary and sufficient condition for the existence of weighted potential, and provides the functional form of the potential function. All proofs are relegated to [Appendix B](#).

Proposition 1. *The game Γ is a weighted potential game if and only if payoffs are quasilinear. In addition, the following functional is a weighted potential of Γ :*

$$V(\alpha, F, \theta) = \int_0^{\bar{\alpha}(\alpha)} u(z, \theta) dz + \int_{\underline{w}}^{\bar{w}} v(\alpha(w), \theta, w) dF(w). \quad (4)$$

The necessity of quasilinear payoffs is brought about by symmetry restrictions imposed by the existence of potential. Roughly speaking, for a single function to reflect payoff differences for all types, (i) the payoff function must exhibit symmetry with respect to the aggregate action and, (ii) the (infinitesimal) contribution to the change in potential of a type switching actions associated with the (infinitesimal) change in the aggregate action must depend only on the size of the switch. Condition (i) leads to separability of payoffs with respect to aggregate action and types, while (ii) implies linearity in own action of the payoff component associated with the aggregate action. For the continuous-action case, externality symmetry translates into equal cross-partial derivatives:

$$\frac{\partial^2 U(a, \bar{a}, \theta, w)}{\partial a \partial \bar{a}} = \frac{\partial^2 U(a', \bar{a}, \theta, w)}{\partial a' \partial \bar{a}} \text{ for all } a, a', \bar{a} \text{ and } w.$$

⁸Product spaces are assumed to be endowed with the product topology.

4 Equilibrium Selection

The characterization of both payoffs and the potential function allows us to establish the equivalence between two commonly used equilibrium selection rules in large games with strategic complementarities (i.e., supermodular games): potential maximization and the global games selection. First, we introduce potential maximization and show that it is associated with finding the NE that maximizes the ex-ante payoffs of an agent with marginal beliefs, i.e., who thinks that types lower than hers do not contribute to the aggregate action. In games with strategic complementarities, marginal beliefs can be interpreted as factoring in miscoordination risk, given that they systematically underestimate the value of the aggregate action. Second, after introducing the global games selection, we show that both selection rules coincide in supermodular games with quasilinear payoffs.

4.1 Potential Maximization

This section introduces the notion of marginal beliefs and conveys two results. [Proposition 2](#) shows that a strategy profile maximizing potential exists and maximizes the ex-ante payoffs of an agent with marginal beliefs. [Proposition 3](#) establishes its uniqueness in supermodular games ([Milgrom and Roberts, 1990](#); [Vives, 1990](#); [Van Zandt and Vives, 2007](#)).

Definition 3 (Marginal beliefs). An agent of type w has *marginal beliefs* with respect to profile α if she believes that the mass of players in the population following strategy α is $1 - F(w)$, and that the remaining players choose $a = 0$. Accordingly, she believes that the aggregate action is given by $\bar{\alpha}(\alpha, [w, \bar{w}])(1 - F(w))$.

Marginal beliefs have two related properties. First, a player thinks that her type is a pivotal type, i.e., she thinks types lower than hers do not contribute to the aggregate action. Second, she underestimates the value of the aggregate action since $\bar{\alpha}(\alpha, [w, \bar{w}]) \leq \bar{\alpha}(\alpha, [\underline{w}, \bar{w}])$.

A strategy profile α_P maximizes the ex-ante expected payoffs of an agent with marginal beliefs if it satisfies

$$\alpha_P \in \arg \max_{\alpha \in \mathcal{A}} \int_{\underline{w}}^{\bar{w}} \left(\alpha(w)u(\bar{\alpha}(\alpha, [w, \bar{w}])(1 - F(w)), \theta) + v(\alpha(w), \theta, w) \right) dF(w). \quad (5)$$

The next result establishes that the set of potential maximizers coincides with the set of profiles that maximize ex-ante payoffs under marginal beliefs.⁹ In addition, it shows that potential maximizers exist and are always NE. To facilitate the exposition, results stating that two strategy profiles are equal are meant to say that they are equal except in a set of (w, θ) with zero Lebesgue measure, unless otherwise noted. Similarly, uniqueness also refers to being unique almost everywhere in $[\underline{w}, \bar{w}] \times \Theta$. Also, since the distribution of types F remains fixed in the subsequent analysis, we denote $V(\alpha, F, \theta)$ by simply $V(\alpha, \theta)$ to ease on notation.

Proposition 2. *If Γ is a weighted potential game then strategy profile α^* maximizes the weighted potential given by (4) if and only if it satisfies (5). In addition, such a profile exists and it is a NE of the game.*

It is worth noting that while marginal beliefs do not necessarily reflect how individual incentives compare across types, in supermodular games such beliefs are coherent with the fact that higher types prefer higher actions.

Assumption 1 (Supermodular game). *Payoffs satisfy*

- (i) ΔU is bounded and Lipschitz continuous.
- (ii) If $a > a'$ then $\Delta U(a, a', \bar{a}, \theta, w)$ is strictly increasing in \bar{a} and also in w . That is, U exhibits strictly increasing differences w.r.t. a and both \bar{a} and w .
- (iii) There exists $K > 0$ such that $U(a, \bar{a}, \theta, w) - U(a', \bar{a}, \theta', w) \geq K(a - a')(\theta - \theta')$ for all $a \geq a'$, all $\theta \geq \theta'$, all \bar{a} and all w .

Increasing differences with respect to aggregate action in part (ii) lead to strategic complementarities. Similarly, part (iii) implies that a higher θ increases the incentives to take higher actions.

Proposition 3. *If Γ is a weighted potential game and satisfies Assumption 1 then there is a unique profile α_P maximizing potential. In addition, α_P is increasing in w and θ .*

⁹The beliefs associated with potential maximization are not unique. In general, if a player of type w has marginal beliefs then there exist permutations of types $r_w : [\underline{w}, \bar{w}] \rightarrow [\underline{w}, \bar{w}]$ satisfying $r_w(w) = w$ such that the agent believes that the aggregate action is given by

$$\bar{\alpha}(\alpha, \{w' : r_w(w') \geq w\}) := \int_{\underline{w}}^{\bar{w}} \alpha(w') dF(w' | r_w(w') \geq w). \quad (6)$$

In what follows, we use $\alpha_P(w, \theta)$ to denote the individual strategies associated with the potential maximizer at θ .¹⁰

The presence of strategic complementarities not only leads to a unique potential maximizer but, as mentioned above, it also imbues it with a natural interpretation in terms of risk: potential maximization favors equilibria that involve less 'miscoordination risk' over those that might be associated with higher average payoffs, given that marginal beliefs underestimate the value of the aggregate action for each type.

We next use the investment game of [Example 1](#) to illustrate how to characterize the potential maximizing equilibrium and discuss the economic implications of maximizing expected payoffs under marginal beliefs. Recall that individual payoffs are given by

$$au(\bar{a}, \theta) - \frac{a^2}{2} \frac{1}{w},$$

which are strictly concave in a . Hence, the optimal investment for an agent of type w , i.e., the best response to aggregate investment \bar{a} , is the solution to the FOC¹¹

$$\alpha(w) = u(\bar{a}, \theta)w. \tag{7}$$

Integrating individual actions across types and recalling that $Ew = 1$ we obtain the equilibrium condition on aggregate investment:

$$\bar{a} = \int_w u(\bar{a}, \theta)w dF(w) = u(\bar{a}, \theta). \tag{8}$$

The solutions \bar{a}^* to this equation represent the NE levels of aggregate investment of the game. Given (7) and (8), we can write equilibrium strategies as $\alpha^*(w) = \bar{a}^*w$.

Next, consider the following s-shaped specification of returns: $u(\bar{a}, \theta) = 2\theta \frac{\bar{a}^2}{\bar{a}^2 + 1}$. Under s-shaped returns, [eq. \(8\)](#) has at most three solutions, zero investment ($\bar{a}^* = 0$) and the real solutions to quadratic equation

$$\bar{a}^{*2} - 2\theta\bar{a}^* + 1 = 0 \quad \Rightarrow \quad \bar{a}^* = \theta \pm (\theta^2 - 1)^{1/2}.$$

¹⁰Given the monotonicity and generic uniqueness of α_P , if there are multiple potential maximizers at θ we can pick $\alpha_P(w, \theta)$ to be the largest of them.

¹¹To keep things simple we assume that the upper bound a_{max} on investment is high enough so that it is not binding for the values of θ that we consider.

That is, the game has multiple NE when $\theta \geq 1$. It is worth noting that the NE exhibiting the largest investment Pareto dominate the lower investment equilibria.¹²

By [Proposition 1](#) the potential function is given by

$$V(\alpha, \theta) = \int_0^{\bar{\alpha}(\alpha)} u(z, \theta) dz - \int_w \frac{\alpha(w)^2}{2} \frac{1}{w} dF(w). \quad (9)$$

Since NE satisfy $\alpha^*(w) = \bar{a}^* w$ and strategy profiles that maximize potential are NE, we can restrict attention to individual strategies of the form $\alpha_{\bar{a}}(w) = \bar{a} w$ and look for the set of \bar{a} that maximize

$$V(\alpha_{\bar{a}}, \theta) = \int_0^{\bar{a}} 2\theta \frac{z^2}{z^2 + 1} dz - \int_w \frac{\bar{a}^2 w^2}{2} \frac{1}{w} dF(w) = 2\theta (\bar{a} - \tan^{-1}(\bar{a})) - \frac{\bar{a}^2}{2}. \quad (10)$$

The left plot in [Figure 1](#) shows that this function has at most two local maximizers, the zero investment equilibrium and the largest solution to [eq. \(8\)](#) whenever it exists.¹³ As the quality of the technology crosses threshold $\hat{\theta} \approx 1.1$, the global maximizer, denoted by \bar{a}_P , switches. This implies that individual investment strategies remain at zero at quality levels below the threshold and discontinuously jump at $\hat{\theta}$, as depicted in the right graph of [Figure 1](#). Specifically, they are given by the cutoff function

$$\alpha_P(w, \theta) = \begin{cases} 0 & \theta < \hat{\theta} \\ \left(\theta + (\theta^2 - 1)^{1/2}\right) w & \theta \geq \hat{\theta}. \end{cases} \quad (11)$$

The zero-investment trap and the discontinuous switch to the high investment equilibrium is due to s-shaped returns, which imply that complementarities are small at low and high investment levels, while being stronger at moderate values of \bar{a} . Maximizing the ex-ante payoffs of an agent with marginal beliefs leads to selecting the no investment equilibrium over the Pareto dominant NE for all $\theta \in [1, \hat{\theta})$. This is driven by the fact that marginal beliefs underestimate investment levels, making the largest investment equilibrium look too risky from an ex-ante perspective when the quality of the technology θ is not very high.

¹²Equilibrium payoffs can be written as $U(\alpha^*(w), \bar{a}^*, \theta, w) = (\bar{a}^* w) \bar{a}^* - \frac{(\bar{a}^* w)^2}{2} \frac{1}{w} = \frac{\bar{a}^*}{2} w$, which are strictly increasing in \bar{a}^* .

¹³The NE with intermediate investment levels is a local minimizer.




Figure 1: Potential maximization (left) and investment strategies (right).

4.2 Global Games Selection

We next describe the Global Games (GG) equilibrium selection rule for supermodular games (Carlsson and van Damme, 1993; Frankel et al., 2003), which is based on introducing incomplete information about parameter θ via idiosyncratic noise. Such noise generates uncertainty about the aggregate action, impairing agents' ability to coordinate via self-fulfilling beliefs, leading to a unique equilibrium. Specifically, each agent gets a signal $s = \theta + \nu\eta$, where $\nu > 0$ is the noise scale, and η is independently distributed according to continuous distribution H_w with full support on $[-1/2, 1/2]$ and density h_w , which is allowed to depend on the agent's type. Agents have a common prior about θ with continuous density ϕ and full support on Θ . We assume that the exact Law of Large Numbers (LLN) applies for each type w , i.e., the mass of agents of type w with signal noise less than ε is given by $H_w(\varepsilon)$. The strategy of a player in the incomplete information game is a mapping from the space of signals S to actions in A . Abusing notation, let $\alpha : [\underline{w}, \bar{w}] \times S \rightarrow A$ denote the strategy profile that assigns action $\alpha(w, s)$ to a player of type w receiving signal s .

The goal of the GG selection is to induce uniqueness of Bayes Nash equilibrium in the incomplete information game and then select an equilibrium of the complete information game as a function of θ by taking the limit as $\nu \rightarrow 0$.

To obtain uniqueness, in addition to Assumption 1, we assume the presence of dominance regions, that is, ranges of parameter values at which all player types have a strictly dominant strategy.

Assumption 2. *There exist $\underline{\theta} > \inf \Theta$ and $\bar{\theta} < \sup \Theta$ such that, for all w and all*

$\bar{a} \in [0, a_{max}]$, if $\theta < \underline{\theta}$ then $\Delta U(a, 0, \bar{a}, \theta, w) < 0$ for all $a > 0$, and if $\theta > \bar{\theta}$ then $\Delta U(a_{max}, a, \bar{a}, \theta, w) > 0$ for all $a < a_{max}$.

The next two propositions establish, respectively, that there is a unique equilibrium (i.e., unique except in a zero measure subset of $[\underline{w}, \bar{w}] \times S$), and that it converges to a NE of Γ as noise vanishes.

Proposition 4. *If Γ satisfies [Assumptions 1 and 2](#) then there exists $\bar{\nu} > 0$ such that for all $\nu < \bar{\nu}$ there is a unique equilibrium in the global game. Moreover, the equilibrium strategy profile is monotone in both s and w .*

The proof is based on standard arguments in the global games literature ([Frankel et al., 2003](#)). First, payoffs exhibiting increasing differences w.r.t. signals and actions implies that the global game is a supermodular game. Thus, the game has both a smallest and a largest equilibrium (Theorem 5 in [Milgrom and Roberts, 1990](#)). Moreover, players follow monotone strategies in these equilibria. Second, we show that under a uniform prior shifting up all strategies w.r.t. signals by the same amount does not affect agents' beliefs about the aggregate action at any given signal, while it does lead to higher expectations about θ . We exploit this translation invariance to prove that, as we move up from the smallest to the largest equilibrium, expected payoffs differences between higher and lower actions go up and thus there can be only one equilibrium. Finally, we show that beliefs (and hence expected payoffs) under a non-uniform prior converge to the beliefs associated with a uniform prior.

Proposition 5. *Let Γ satisfy [Assumptions 1 and 2](#). There exists a strategy profile α_G such that, for any sequence α^ν of equilibria in the global game indexed by $\nu \rightarrow 0$, $\lim_{\nu \rightarrow 0} \alpha^\nu(w, s) = \alpha_G(w, s)$ for almost all w, s . Moreover, $\alpha_G(\theta, \cdot)$ is a NE of Γ for almost all $\theta \in \Theta$.*

A desirable property of the GG selection is to be invariant to different noise distributions H_w , given that the goal is to pin down equilibrium in the limit and, hence, the introduction of noise is just a convenient technical device to introduce miscoordination risk. In such a case, the GG selection is said to be *noise-independent*. As we show next, this is indeed the case under quasilinear payoffs, given its equivalence to potential maximization.

4.3 Equivalence Result

[Theorem 1](#) establishes below that the limit equilibrium in the global game coincides with the potential maximizing strategy profile in the complete information game. We obtain the result by identifying a key property of beliefs about the aggregate action in the global game, which we call the *Generalized Laplacian Property* (GLP). The GLP links the (weighted) average belief to the uniform distribution. We use the GLP to show that, under quasilinear payoffs, the change in expected payoffs of a type that switches actions coincides with the change in potential as noise vanishes. Since this is a key step in the proof of equivalence, we use [Example 1](#) to introduce the GLP and provide some intuition before formally stating the theorem.

Recall that, in the investment game, the potential maximizing profile $\alpha_P(w, \theta)$ given by (11) implies that firms follow a cutoff rule, i.e., they choose zero investment if $\theta < \hat{\theta}$ and the largest NE investment if $\theta \geq \hat{\theta}$ (see [Figure 1](#)). Accordingly, for the limit equilibrium in the global game to coincide with $\alpha_P(w, \theta)$, a firm with type w needs to be indifferent between choosing $\alpha_1(w) = 0$ and $\alpha_2(w) = \alpha_P(w, \hat{\theta})$ when it receives signal $s = \hat{\theta}$. That is, the expected payoff difference between the two investment levels conditional on $s = \hat{\theta}$ must be zero. We partially illustrate how this is the case by showing that the average expected payoff difference coincides with the difference in potential between strategy profiles α_1 and α_2 , which is zero since both profiles maximize potential at $\hat{\theta}$.

To do so, assume that in the global game a firm of type w chooses $\alpha_1(w)$ if its signal is below some cutoff $\kappa(w)$ and $\alpha_2(w)$ if its signal is above $\kappa(w)$. Furthermore, assume that firms have a uniform prior over Θ and that their cutoffs $\kappa(\cdot)$ are within noise range ν of each other. Notice that, when ν is very small, θ is very close to the value of the firm's signal, and expected payoffs conditional on signal $s = \kappa(w)$ can be approximated by

$$E(U(a, \bar{a}, \theta, w) | s = \kappa(w)) \approx \int_z au(z, \kappa(w)) dG_w(z) - \frac{a^2}{2} \frac{1}{w},$$

where $G_w(z) = Pr(\bar{a} \leq z | s = \kappa(w), w)$ denotes the cdf of aggregate investment conditional on signal $s = \kappa(w)$ and on type w when players follow the above cutoff strategies. Hence, the difference in expected payoffs across the two actions can be

expressed as

$$0 \approx \int_z (\alpha_2(w) - \alpha_1(w)) u(z, \kappa(w)) dG_w(z) - \frac{\alpha_2(w)^2 - \alpha_1(w)^2}{2} \frac{1}{w}.$$

Solving these indifference conditions requires pinning down G_w , which depends on the distribution of noise and signal cutoffs κ . However, under quasilinear payoffs and a uniform prior, we can circumvent this problem by focusing on *average* instead of individual indifference conditions and using the GLP to replace average beliefs. The next lemma presents the GLP under a uniform prior, which states that the weighted average of beliefs G_w is given by the uniform distribution. The weights represent the contribution of type w to the aggregate action when it switches actions, i.e., they are given by $(\alpha_2(w) - \alpha_1(w))f(w)$.

Lemma 1 (Generalized Laplacian Property). *Fix $\alpha_1, \alpha_2 \in \mathcal{A}$ satisfying $\alpha_1(w) \leq \alpha_2(w)$ for all w . If players have a uniform prior then, for any signal cutoff function $\kappa : [\underline{w}, \bar{w}] \rightarrow [\inf \Theta + \nu/2, \sup \Theta - \nu/2]$ such that each type w chooses $\alpha_1(w)$ if $s < \kappa(w)$ and $\alpha_2(w)$ if $s \geq \kappa(w)$ and for all $z \in [\bar{\alpha}(\alpha_1), \bar{\alpha}(\alpha_2)]$, we have that¹⁴*

$$\int_{\underline{w}}^{\bar{w}} (\alpha_2(w) - \alpha_1(w)) G_w(z) dF(w) = z - \bar{\alpha}(\alpha_1). \quad (12)$$

Averaging indifference conditions across firms we obtain

$$0 \approx \int_w \int_z (\alpha_2(w) - \alpha_1(w)) u(z, \kappa(w)) dG_w(z) dF(w) - \int_w \frac{\alpha_2(w)^2 - \alpha_1(w)^2}{2} \frac{1}{w} dF(w).$$

In addition, as ν goes to zero, since firms' signal cutoffs are within the noise range of each other, $u(z, \kappa(w))$ can be approximated by $u(z, k)$, where k is the limit cutoff to which all $\kappa(w)$ converge. Hence, the first integral in the above expression satisfies

$$\int_z u(z, k) \int_w (\alpha_2(w) - \alpha_1(w)) dG_w(z) dF(w) = \int_{\bar{\alpha}(\alpha_1)}^{\bar{\alpha}(\alpha_2)} u(z, k) dz,$$

where the equality comes from applying (12) to replace the average belief while the

¹⁴To see how (12) implies that \bar{a} is uniformly distributed in $[\bar{\alpha}(\alpha_1), \bar{\alpha}(\alpha_2)]$ note that we can divide both sides by $\bar{\alpha}(\alpha_2) - \bar{\alpha}(\alpha_1)$ so that the RHS is equal to uniform cdf $\frac{z - \bar{\alpha}(\alpha_1)}{\bar{\alpha}(\alpha_2) - \bar{\alpha}(\alpha_1)}$.

limits of integration reflect the range of feasible aggregate investment levels given the cutoff strategies used by firms. This implies that cutoff k must satisfy

$$0 \approx \int_{\bar{\alpha}(\alpha_1)}^{\bar{\alpha}(\alpha_2)} u(z, k) dz - \int_w \frac{\alpha_2(w)^2 - \alpha_1(w)^2}{2} \frac{1}{w} dF(w).$$

But notice that, when $k = \hat{\theta}$, the RHS is the difference in potential between NE profiles α_2 and α_1 at quality $\hat{\theta}$, which is zero since both profiles maximize potential.

The relationship between expected payoff differences and changes in potential partially illustrated by this example can be established for each type individually by applying the GLP to any subset of types, effectively linking changes in expected payoffs following a deviation from the potential maximizing equilibrium to changes in potential. This leads to the following equivalence result.

Theorem 1. *If Γ is a weighted potential game and satisfies [Assumptions 1 and 2](#) then α_G is equal to α_P for any prior with continuous density and full support in Θ , implying that α_G is noise-independent.*

The equivalence result provides a tractable characterization of the GG selection as well as an economic interpretation in terms of maximizing ex-ante payoffs under marginal beliefs. In addition, it identifies quasilinearity as a sufficient, easy-to-check condition for noise independence in aggregative games.

The proof works as follows. It first shows for the uniform-prior case that, for any subset of types W , the expected average payoff gain across types in W from deviating from α_P to any profile α using a common signal threshold $s = \theta$ is equal to the change in potential. This is done by applying a version of the GLP showing that the average belief about the aggregate action of types in W is the uniform distribution ([Lemma 3 in Appendix A](#)). Since we can choose W to be a small neighborhood of any type w and potential maximization implies that the change in potential must be negative, a continuity argument implies that, for almost all types, the deviation from α_P to α is not profitable, i.e., that the potential maximizing equilibrium profile coincides with the limit equilibrium in the global game. The proof then uses a limit version of the GLP for non-uniform priors to extend the equivalence result beyond the uniform-prior case ([Lemma 5 in Appendix A](#)).¹⁵

¹⁵The limit version of the GLP could be used to relax the assumption of additive noise. For

The GLP holds irrespective of the payoff structure and is the product of combining two key ingredients, namely, monotone cutoff strategies and uniform prior with additive noise. Cutoff strategies associate the aggregate action with signal quantiles: the aggregate action of any given type w is given by the proportion of agents with signals higher than the type's cutoff, weighted by the difference in actions ($\alpha_2(w) - \alpha_1(w)$). The uniform prior and additive noise lead to signals $s = \theta + \nu\eta$ being uniformly distributed and independent of types, as long as signals lie in $[\inf \Theta + \nu/2, \sup \Theta - \nu/2]$. This is because, as shown by [Lemma 4](#) in [Appendix A](#), the sum of two independent random variables, one of them uniformly distributed and with a larger support than the other, has a constant density except at the tails. Such a statistical fact explains the connection between average beliefs and the uniform distribution and also their invariance with respect to H_w , since only the prior determines the distribution of signals (except for signals very close to the boundaries of their support).

The above example also illustrates the key role that quasilinearity plays in the characterization of the global games selection and, particularly, in its independence of the noise structure. If the payoff impact of the aggregate action is not linear in own action or it is asymmetric across types then we cannot apply the GLP to replace beliefs, either because belief weights differ from the difference in actions or because the average belief cannot be separated from payoffs. Hence, solving for equilibrium would require the use of individual beliefs, which depend in non-trivial ways on the distribution of noise (see [Serrano-Padial \(2018\)](#) for an example in which the GG selection depends on H_w when payoffs are not quasilinear).

5 Robust Comparative Statics

This section identifies conditions under which changes in parameters (θ) or heterogeneity (F) makes the set of NE aggregate actions to go up. By exploiting the quasilinear payoff structure, we are able to provide conditions on average payoffs.

instance, any signals of the form $s = d(\theta, \eta; \nu)$ with d strictly increasing for which there exists a monotone transformation $q(s) = q_1(\theta) + \nu q_2(\eta)$ would lead to the same limit equilibrium. These include multiplicative noise, i.e., $s = \theta\eta^\nu$, as well as exponential noise ($s = \theta\eta^\nu$), with noise support defined to ensure that signals are well-defined and monotone in θ and η . The reason is that we can redefine signals as $q(s)$, the common value parameter as $q_1(\theta)$ and noise as $q_2(\eta)$ so that the associated distributions satisfy all the assumptions, thus leading to the same equilibrium selection.

To do so, we first characterize the set of NE aggregate actions as fixed points of a nested maximization problem.

Let $\mathcal{A}_{\bar{a}} = \{\alpha \in \mathcal{A} : \bar{\alpha}(\alpha) = \bar{a}\}$ be the set of strategy profiles with aggregate action \bar{a} . In addition, denote by $B(\bar{a}, \theta)$ the maximum average value of the idiosyncratic payoff component $v(a, \theta, w)$ when strategy profiles are restricted to $\mathcal{A}_{\bar{a}}$, that is,

$$B(\bar{a}, \theta) := \max_{\alpha \in \mathcal{A}_{\bar{a}}} \int_w v(a, \theta, w) dF(w). \quad (13)$$

Lemma 2. *If payoffs are quasilinear then \bar{a}^* is the aggregate action in some NE if and only if*

$$\bar{a}^* \in \arg \max_{\bar{a} \in [0, a_{max}]} (\bar{a}u(\bar{a}^*, \theta) + B(\bar{a}, \theta)). \quad (14)$$

This result allows us to establish conditions on $u(\bar{a}, \theta)$ and $B(\bar{a}, \theta)$ for the set of NE aggregate actions to go up after an increase in θ ([Theorems 2](#) and [3](#)) or after a change in the type distribution F ([Theorems 4](#) and [5](#)).

Theorem 2 (Robust Comparative Statics I). *Let payoffs be quasilinear and Θ' a compact subset of Θ . If $u(\bar{a}, \theta)$ is increasing in $\theta \in \Theta'$ for all $\bar{a} \in [0, a_{max}]$ and $B(\bar{a}, \theta)$ exhibits strictly increasing differences in $[0, a_{max}] \times \Theta'$ then the smallest and largest NE aggregate actions are increasing in Θ' .*

[Theorem 2](#) can be further generalized by assuming that $\bar{a}u(\bar{a}', \theta) + B(\bar{a}, \theta)$ is single-crossing in \bar{a}, θ for all \bar{a}' ([Milgrom and Shannon, 1994](#)). However, imposing separate conditions on u and B allows for a simpler derivation of direct restrictions on payoffs. Nonetheless, the conditions on $B(\bar{a}, \theta)$ can be hard to interpret since they involve the value function of a constrained maximization problem. The next result provides sufficient conditions for the smallest and a largest NE to go up with θ in the continuous-action case. Let $\text{int}(X)$ denote the interior of a set X . In particular, $\text{int}(\mathcal{A})$ is the set of measurable profiles such that $\alpha(w) \in (0, a_{max})$ for all w .

Theorem 3. *Let $A = [0, a_{max}]$ and $\Theta' \subseteq \Theta$ be a closed interval of parameter values θ . Let $u(\bar{a}, \theta)$ be increasing in $\theta \in \Theta'$ for all $\bar{a} \in [0, a_{max}]$. If the smallest and a largest NE are in $\text{int}(\mathcal{A})$ for all $\theta \in \Theta'$ and, for any $\alpha \in \text{int}(\mathcal{A})$ and any $\theta \in \text{int}(\Theta')$,*

$$\int_{\underline{w}}^{\bar{w}} \frac{\partial^2 v(a, \theta, w)}{\partial a \partial \theta} \Big|_{a=\alpha(w)} dF(w) > 0, \quad (15)$$

then the smallest and a largest NE aggregate actions are increasing in Θ' .

Condition (15) is straightforward to check and it requires the idiosyncratic payoff component to exhibit *average* increasing differences in $\text{int}(\mathcal{A}) \times \text{int}(\Theta')$. For instance, in the context of [Example 2](#) this amounts to the expected marginal benefit from consumption being increasing in θ . To see why, recall that in the example payoffs exhibit negative externalities and are given by

$$U(a, \bar{a}, \theta, w) = b(a, \theta, w) - c(\bar{a}, \theta)a,$$

where b is strictly concave in a and c is increasing in \bar{a} . Concavity of b implies that utility is strictly concave. Hence, as long as the marginal benefit satisfies $\frac{\partial b(a, \theta, w)}{\partial a} \Big|_{a=0} > c(\bar{a}, \theta)$ and a_{max} is large enough, the optimal consumption levels will lie in the interior of $[0, a_{max}]$. Hence, condition (15) translates into

$$\int_{\underline{w}}^{\bar{w}} \frac{\partial^2 b(a, \theta, w)}{\partial a \partial \theta} \Big|_{a=\alpha(w)} dF(w) = \frac{\partial}{\partial \theta} E \left(\frac{\partial b(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} \right) > 0.$$

This condition is significantly weaker than imposing monotonicity restrictions pointwise at the type level ([Milgrom and Shannon, 1994](#); [Acemoglu and Jensen, 2010](#)). For instance, it is straightforward to check that if payoffs exhibit increasing differences for all types then they satisfy the conditions in [Theorem 3](#). In contrast, [Theorem 3](#) allows for individual payoffs of a subset of types to exhibit strict *decreasing* differences in a, θ .¹⁶ In addition, the theorem presents conditions on payoffs, which are primitives of the game, making them easier to check than restrictions on average best responses ([Camacho et al., 2018](#)).

Theorem 4 (Robust Comparative Statics II). *Let $B(\bar{a}, \theta)$ and $\hat{B}(\bar{a}, \theta)$ be the value functions defined by (13) associated with type distributions F and \hat{F} , respectively. If*

$$B(\bar{a}, \theta) - B(\bar{a}', \theta) \geq \hat{B}(\bar{a}, \theta) - \hat{B}(\bar{a}', \theta)$$

for any $\bar{a}, \bar{a}' \in [0, a_{max}]$ such that $\bar{a} > \bar{a}'$, then the smallest and largest NE aggregate actions are higher under F than under \hat{F} .

¹⁶In [Example 2](#) payoffs U exhibit decreasing differences if $\frac{\partial^2 U}{\partial a \partial \theta} \leq 0$, that is, if $\frac{\partial^2 b}{\partial a \partial \theta} < \frac{\partial c}{\partial \theta}$.

The next theorem presents sufficient conditions that are easy to verify for games with continuous actions.

Theorem 5. *Let $A = [0, a_{max}]$ and F, \hat{F} be two type distributions. If the smallest and largest NE are in $\text{int}(A)$ for both F and \hat{F} , and, for any $\alpha \in \text{int}(A)$,*

$$\int_{\underline{w}}^{\bar{w}} \frac{\partial v(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} dF(w) > \int_{\underline{w}}^{\bar{w}} \frac{\partial v(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} d\hat{F}(w), \quad (16)$$

then the smallest and largest NE aggregate actions are higher under F than under \hat{F} .

In [Example 2](#) condition (16) boils down to

$$\int_{\underline{w}}^{\bar{w}} \frac{\partial b(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} dF(w) \geq \int_{\underline{w}}^{\bar{w}} \frac{\partial b(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} d\hat{F}(w).$$

This is the case, for instance, if $\frac{\partial b}{\partial a}$ is increasing in w and F first-order stochastically dominates \hat{F} , or if $\frac{\partial b}{\partial a}$ is concave in w and \hat{F} is a mean-preserving spread of F .

We finish by mentioning that, although beyond the scope of the paper, the above characterization of the set of NE aggregate actions ([Lemma 2](#)) could be used to explore whether there exist average conditions on payoffs that guarantee uniqueness of the equilibrium aggregate action, instead of relying on pointwise conditions.¹⁷

6 Related Literature

Our discussion of the related literature narrowly focuses on the three more closely related areas, namely, large potential games, heterogeneous global games, and comparative statics in aggregative games.

[Monderer and Shapley \(1996\)](#) introduced various definitions of potential in finite games. Among other properties, potential maximizers have been shown to be evolutionary stable. In the context of large games, [Sandholm \(2009\)](#) defined potential in population games with finite actions and finite types, and showed that the existence of potential in these games is linked to payoffs exhibiting externality

¹⁷[Cheung and Lahkar \(2018\)](#) and [Lahkar \(2017\)](#) study equilibrium existence in potential aggregative games with a homogeneous population.

symmetry.¹⁸ Our notion of potential extends the definition of potential to allow for continuous actions and types, and we show why the existence of potential implies quasilinear payoffs by deriving symmetry conditions for payoffs that depend on the aggregate action. We also identify the potential function for this class of aggregative games, leading to a tractable characterization of potential maximizing equilibria, and provide economic content behind potential maximization, which has been an open question in the literature.

Frankel et al. (2003) proposed the global games selection for games with heterogeneous payoffs and established uniqueness of the selected equilibrium in both finite- and continuum-player games with finite types. Focusing on binary-action games, Sakovics and Steiner (2012) identified a key property of average beliefs that they used to characterize the global games selection. Drozd and Serrano-Padial (2018) extended such characterization to binary action games with asymmetric equilibria. Our results build upon their work by generalizing the belief property beyond binary actions and by characterizing the global games selection using the potential function. The connection between potential maximization and the global games selection can be established for finite supermodular games by combining the results of Ui (2001) and Morris and Ui (2005), who respectively show that maximizers of potential and the more general notion of monotone potential are robust to incomplete information in the sense of Kajii and Morris (1997),¹⁹ with those of Basteck et al. (2013), who show that the global games selection picks the robust equilibrium whenever it exists. In addition to directly proving this connection in large aggregative games, since the potential maximizing equilibrium is unrelated to the distributional assumptions about noise used in the global game, we provide an easy-to-check sufficient condition, quasilinearity, for the global games selection to be noise independent.

Finally, the paper contributes to the recent literature of aggregate comparative statics (Acemoglu and Jensen, 2010, 2015; Camacho et al., 2018), which focuses on comparative statics on aggregate behavior instead of on individual choices (Topkis, 1979; Milgrom and Roberts, 1990; Vives, 1990; Milgrom and Shannon, 1994). Acemoglu and Jensen (2010, 2015) find monotonicity conditions on individual best

¹⁸Hofbauer and Sandholm (2007) and Zusai (2018) prove the evolutionary stability of local maximizers of potential in population games with finite actions and player types.

¹⁹Potential maximization implies monotone potential maximization. Oyama and Takahashi (2020) prove that monotone potential maximization is necessary and sufficient for robustness in binary-action finite games.

responses for the smallest and largest equilibrium aggregate actions to be monotone in the model parameters. [Camacho et al. \(2018\)](#) further relax these restrictions by pinning down monotonicity conditions on average best responses. We expand their contributions by identifying direct restrictions on average payoffs.

Appendix A The Generalized Laplacian Property

This section presents the full version of the GLP for uniform prior and the limit version of the GLP for non-uniform priors.

Lemma 3 (Generalized Laplacian Property). *Assume agents have a uniform prior. Fix any measurable subset of types W , any $\alpha_1, \alpha_2 \in \mathcal{A}$ with $\alpha_1(w) \leq \alpha_2(w)$ for all $w \in W$, and any cutoff function $\kappa : [\underline{w}, \bar{w}] \rightarrow [\inf \Theta + \nu/2, \sup \Theta - \nu/2]$ such that $\alpha(w, s) = \alpha_1(w)$ if $s < \kappa(w)$ and $\alpha(w, s) = \alpha_2(w)$ if $s \geq \kappa(w)$ for all $w \in W$. Then, for all $z \in [\bar{\alpha}(\alpha_1, W), \bar{\alpha}(\alpha_2, W)]$,*

$$\int_W (\alpha_2(w) - \alpha_1(w)) G_w(z | \kappa; \alpha, W) dF(w | w \in W) = z - \bar{\alpha}(\alpha_1, W), \quad (17)$$

where $G_w(z | \kappa; \alpha, W) := Pr(\bar{\alpha}(\alpha, W) < z) | s = \kappa(w), w$.

To prove [Lemma 3](#) we make use of the following property of the sum of two independent random variables.

Lemma 4. *Let x, y be two independent random variables such that x is uniformly distributed in $[\underline{x}, \bar{x}]$ and y has a density f_y with support $[\underline{y}, \bar{y}]$. If $\bar{y} - \underline{y} < \bar{x} - \underline{x}$ then the sum $z = x + y$ has a constant density in $[\underline{x} + \bar{y}, \bar{x} + \underline{y}]$. Specifically,*

$$f_z(z) = \begin{cases} \frac{1}{\bar{x} - \underline{x}} F_y(z - \underline{x}) & z < \underline{x} + \bar{y} \\ \frac{1}{\bar{x} - \underline{x}} & z \in [\underline{x} + \bar{y}, \bar{x} + \underline{y}] \\ \frac{1}{\bar{x} - \underline{x}} (1 - F_y(z - \bar{x})) & z > \bar{x} + \underline{y}. \end{cases} \quad (18)$$

Proof of Lemma 4. Note that, since $x = z - y$ for any given z , we must have that $x \in [\max\{\underline{x}, z - \bar{y}\}, \min\{\bar{x}, z - \underline{y}\}]$. The joint density of z and x is given by $f_z(z|x)f_x(x)$. In addition, $f_z(z|x) = f_y(z - x|x) = f_y(z - x)$ since y is independent of x . Hence, the density of z satisfies $f_z(z) = \int_{\underline{x}}^{\bar{x}} f_z(z|x)f_x(x)dx = \int_{\max\{\underline{x}, z - \bar{y}\}}^{\min\{\bar{x}, z - \underline{y}\}} f_y(z - x) \frac{1}{\bar{x} - \underline{x}} dx$, leading to the following expression:

$$f_z(z) = \frac{1}{\bar{x} - \underline{x}} (F_y(z - \max\{\underline{x}, z - \bar{y}\}) - F_y(z - \min\{\bar{x}, z - \underline{y}\})),$$

which yields (18) by plugging the values of $\max\{\underline{x}, z - \bar{y}\}$ and $\min\{\bar{x}, z - \underline{y}\}$ for the following three cases. First, if $z < \underline{x} + \bar{y}$ then $\max\{\underline{x}, z - \bar{y}\} = \underline{x}$ and $\min\{\bar{x}, z - \underline{y}\} = z - \underline{y}$. Second, if $z \in [\underline{x} + \bar{y}, \bar{x} + \underline{y}]$ then $\max\{\underline{x}, z - \bar{y}\} = z - \bar{y}$ and $\min\{\bar{x}, z - \underline{y}\} = z - \underline{y}$. Finally, if $z > \bar{x} + \underline{y}$ then $\max\{\underline{x}, z - \bar{y}\} = z - \bar{y}$ and $\min\{\bar{x}, z - \underline{y}\} = \bar{x}$. \square

Proof of Lemma 3. The proof consists of two parts. The first shows that when agents follow cutoff strategy κ the aggregate action coincides with the aggregate action in a game where the set of available actions is normalized to be $\{0, 1\}$ and the type distribution is weighted by the difference $\alpha_2(w) - \alpha_1(w)$. The second part shows that, in the normalized game, the average belief conditional on $s = k(w)$ about the aggregate action of types in W is uniformly distributed in $[0, 1]$.

Abusing notation, for any $\theta \in \Theta$, let $\bar{\alpha}(\alpha, W, \theta)$ denote the aggregate action under the strategy $\alpha(w, s) = \alpha_1(w)$ if $s < \kappa(w)$ and $\alpha(w, s) = \alpha_2(w)$ otherwise. By the exact LLN, the fraction of agents of type w that receive a signal below cutoff $\kappa(w)$ is given by $1 - H_w\left(\frac{k(w) - \theta}{\nu}\right)$. Accordingly, the aggregate action associated with types in W when they follow cutoff strategy κ is given by

$$\bar{\alpha}(\alpha, W, \theta) = \bar{\alpha}(\alpha_1, W) + \int_W (\alpha_2(w) - \alpha_1(w)) \left(1 - H_w\left(\frac{k(w) - \theta}{\nu}\right)\right) f(w|w \in W) dw.$$

Define the density function $\hat{f}(w|w \in W) = \frac{\alpha_2(w) - \alpha_1(w)}{\bar{\alpha}(\alpha_2, W) - \bar{\alpha}(\alpha_1, W)} f(w|w \in W)$. Note that \hat{f} is well-defined since $\bar{\alpha}(\alpha_i, W) = \int_W \alpha_i(w) f(w|w \in W) dw$ for $i = 1, 2$. Let \hat{F} be the corresponding cdf.

We can express the aggregate action in W as

$$\bar{\alpha}(\alpha, W, \theta) = \bar{\alpha}(\alpha_1, W) + (\bar{\alpha}(\alpha_2, W) - \bar{\alpha}(\alpha_1, W)) y(\kappa, W, \theta), \quad (19)$$

where $y(\kappa, W, \theta) = \int_W \left(1 - H_w\left(\frac{k(w) - \theta}{\nu}\right)\right) \hat{f}(w|w \in W) dw$ represents the mass of agents in W with signals $s \geq \kappa(w)$. Accordingly, we have that

$$\begin{aligned} & \int_W (\alpha_2(w) - \alpha_1(w)) G_w(z|\kappa; \alpha, W) dF(w|w \in W) \\ &= \int_W Pr\left(y(\kappa, W, \theta) < \frac{z - \bar{\alpha}(\alpha_1, W)}{\bar{\alpha}(\alpha_2, W) - \bar{\alpha}(\alpha_1, W)} \mid s = \kappa(w), w\right) d\hat{F}(w|w \in W) \quad (20) \end{aligned}$$

Since $\frac{z - \bar{\alpha}(\alpha_1, W)}{\bar{\alpha}(\alpha_2, W) - \bar{\alpha}(\alpha_1, W)} \in [0, 1]$ when $z \in [\bar{\alpha}(\alpha_1, W), \bar{\alpha}(\alpha_2, W)]$, to prove (17) it suffices to show that

$$\int_W Pr(y(\kappa, W, \theta) < z \mid s = \kappa(w), w) d\hat{F}(w|w \in W) = z \text{ for all } z \in [0, 1]. \quad (21)$$

To do so, consider the normalized game $\hat{\Gamma} = \{\hat{F}, \{0, 1\}, \theta, U\}$. If agents in the global game version of $\hat{\Gamma}$ follow strategy $\alpha(w, s) = 0$ if $s < \kappa(w)$ and $\alpha(w, s) = 1$ otherwise, then the aggregate action in W is given by $y(\kappa, W, \theta)$ for all θ .

To prove (21), define ‘virtual signals’ $\tilde{s} = s - \kappa(w)$ for all $w \in W$, which exhibit a common cutoff $\tilde{\kappa} = 0$. Let the ‘extended type’ of a player be the tuple (s, w) .

Since θ is uniformly distributed in $[\inf \Theta, \sup \Theta]$ and $\nu\eta$ is independent of θ with support $[-\nu/2, \nu/2]$, by Lemma 4, signals in $[\inf \Theta + \nu/2, \sup \Theta - \nu/2]$ have constant density $\frac{1}{\sup \Theta - \inf \Theta}$ independent of w . Accordingly, the density associated with extended type $(k(w), w)$, conditional on $\tilde{s} = 0$ and on $w \in W$, is given by

$$Pr(\kappa(w), w | \tilde{s} = 0, W) = \frac{Pr(k(w), w | W)}{Pr(\tilde{s} = 0 | W)} = \frac{\frac{1}{\sup \Theta - \inf \Theta} \hat{f}(w | W)}{\frac{1}{\sup \Theta - \inf \Theta}} = \frac{\hat{f}(w)}{\int_W \hat{f}(w) dw}. \quad (22)$$

where $Pr(s, w | \cdot)$ denotes the conditional probability density of extended type (s, w) .

Next, we show that $y(\kappa, W, \theta)$ is uniformly distributed conditional on $\tilde{s} = 0$, i.e.,

$$Pr(y(\kappa, W, \theta) < z | \tilde{s} = 0, W) = z. \quad (23)$$

First note that the virtual noise $\tilde{\eta} = (\tilde{s} - \theta)/\nu$ follows the mixture distribution $\left\{ H_w \left(\tilde{\eta} + \frac{\kappa(w)}{\nu} \right), \frac{\hat{f}(w)}{\int_W \hat{f}(w) dw} \right\}_W$. This implies that the virtual noise belongs to type w with probability $\frac{\hat{f}(w)}{\int_W \hat{f}(w) dw}$. In addition, its distribution conditional on type w is given by the noise distribution evaluated at $\eta = \tilde{\eta} + \kappa(w)/\nu$. But note that the mixture distribution does not depend on θ so the random variable $\tilde{\eta}$ is i.i.d. across agents and independent of θ .

Let \hat{H} be the cdf of $\tilde{\eta}$ and define $\hat{H}^{-1}(z) = \inf\{\tilde{\eta} : \hat{H}(\tilde{\eta}) = z\}$. Given the definition of virtual noise, the aggregate action in subset W is given by the fraction of agents in W whose virtual signal is greater than zero, i.e., by one minus the cdf of the virtual noise \hat{H} evaluated at $-\theta/\nu$. This yields expression (23) given that

$$\begin{aligned} Pr(y(\kappa, W, \theta) < z | \tilde{s} = 0, W) &= Pr(1 - \hat{H}(-\theta/\nu) < z | \tilde{s} = 0, W) = Pr(1 - \hat{H}(\tilde{\eta}) < z) \\ &= Pr(\tilde{\eta} > \hat{H}^{-1}(1 - z)) = 1 - \hat{H}(\hat{H}^{-1}(1 - z)) = z. \end{aligned}$$

Combining (22) and (23) we obtain (21), since

$$Pr(y(\cdot) < z | \tilde{s} = 0, W) = \int Pr(y(\cdot) < z | s = \kappa(w), w) Pr(s = \kappa(w), w | \tilde{s} = 0, W) dw. \quad \square$$

Lemma 4 reveals the key role that the uniform prior and additive noise play by inducing a uniform distribution of signals. Nonetheless, a version of the GLP for non-uniform priors approximately holds when noise levels are very small, in which

the weights on individual beliefs depend on the prior.

Lemma 5 (Generalized Laplacian Property for Non-uniform Prior). *Assume agents have a common prior with continuous density ϕ that has full support on Θ . Fix any measurable subset of types W , any $\alpha_1, \alpha_2 \in \mathcal{A}$ with $\alpha_1(w) \leq \alpha_2(w)$ for all $w \in W$, and any cutoff function $\kappa : [\underline{w}, \bar{w}] \rightarrow [\inf \Theta + \nu/2, \sup \Theta - \nu/2]$ such that $\alpha(w, s) = \alpha_1(w)$ if $s < \kappa(w)$ and $\alpha(w, s) = \alpha_2(w)$ if $s \geq \kappa(w)$ for all $w \in W$. Then,*

$$\lim_{\nu \rightarrow 0} \int_W (\alpha_2(w) - \alpha_1(w)) G_w(z|\kappa; \alpha, W) \frac{\phi(\kappa(w))f(w)}{\int_W \phi(\kappa(w))f(w)dw} dw = z - \bar{\alpha}(\alpha_1, W) \quad (24)$$

for all $z \in [\bar{\alpha}(\alpha_1, W), \bar{\alpha}(\alpha_2, W)]$, and the convergence as $\nu \rightarrow 0$ is uniform.

Proof of Lemma 5. The proof adapts the steps of the proof of Lemma 3 to the case of a non-uniform prior. Specifically, we first show that the joint density of extended types (s, w) conditional on $\tilde{s} = 0$ and $w \in W$ uniformly converges to

$$\lim_{\nu \rightarrow 0} Pr(\kappa(w), w | \tilde{s} = 0, W) = \frac{\phi(\kappa(w))\hat{f}(w)}{\int_W \phi(\kappa(w))\hat{f}(w)dw}, \quad (25)$$

Given this, we show that $y(\kappa, W, \theta)$ is uniformly distributed in $[0, 1]$ conditional on $\tilde{s} = 0$ so that condition (23) holds in the limit. Accordingly, combining (25) and (23) we obtain (24).

To prove (25) note that the joint density of (s, w, θ) is now given by

$$Pr(s, w, \theta | W) = Pr(s|w, \theta)Pr(w|W)Pr(\theta) = \left(h_w \left(\frac{s - \theta}{\nu} \right) \frac{1}{\nu} \right) \left(\frac{\hat{f}(w)}{\int_W \hat{f}(w)dw} \right) \phi(\theta).$$

We obtain the marginal density of (s, w) by integrating the above expression, which leads to, after applying the change of variable $\theta' = \frac{s - \theta}{\nu}$,

$$\begin{aligned} Pr(s, w | W) &= \int_{s-\nu/2}^{s+\nu/2} h_w \left(\frac{s - \theta}{\nu} \right) \frac{1}{\nu} \frac{\hat{f}(w)}{\int_W \hat{f}(w)dw} \phi(\theta) d\theta \\ &= \int_{-1/2}^{1/2} h_w(\theta') \frac{\hat{f}(w)}{\int_W \hat{f}(w)dw} \phi(s - \nu\theta') d\theta' \rightarrow \frac{\hat{f}(w)}{\int_W \hat{f}(w)dw} \phi(s) \text{ as } \nu \rightarrow 0. \end{aligned}$$

The limit is continuous, so the convergence of distribution functions is uniform. We obtain condition (25) by taking the limit as $\nu \rightarrow 0$,

$$Pr(s = \tilde{s} + \kappa(w) | W) = \int_W Pr(\tilde{s} + \kappa(w), w | W) dw \rightarrow \frac{\int_W \phi(\tilde{s} + \kappa(w))\hat{f}(w)dw}{\int_W \hat{f}(w)dw}.$$

To show that (23) holds in the limit notice that the distribution of the virtual noise converges to the mixture distribution $\left\{ H_w \left(\tilde{\eta} + \frac{\kappa(w)}{\nu} \right), \frac{\phi(\kappa(w))\hat{f}(w)}{\int_W \phi(\kappa(w))\hat{f}(w)dw} \right\}_W$, which does not depend on θ so $\tilde{\eta}$ is i.i.d. across agents and independent of θ . Hence, the argument in the last part of the proof of Lemma 3 applies. \square

Appendix B Omitted Proofs

B.1 Proofs of Results in Section 3

Proof of Proposition 1. The “if” part of the proof is based on the following steps. First, we show that condition (3) implies that payoffs satisfy *externality symmetry*: the infinitesimal change in the payoff differences of type w when type w' switches actions is the same as the change in payoff differences of type w' when type w switches actions. Second, we show that externality symmetry implies quasilinearity.

For the “only if” part, we construct a potential function for quasilinear payoffs that satisfies condition (3).

“If” part: Assume that there exists functional V satisfying condition (3) and focus on how the infinitesimal change in V due to a switch of type w from a to a' changes when type w' switches from a'' to a''' . Let strategy profiles $\alpha, \alpha', \alpha''$ and α''' satisfy $\alpha(w) = \alpha''(w) = a$, $\alpha'(w) = \alpha'''(w) = a'$, $\alpha(w') = \alpha'(w') = a''$, $\alpha''(w') = \alpha'''(w') = a'''$ and $\alpha(w'') = \alpha'(w'') = \alpha''(w'') = \alpha'''(w'')$ for all $w'' \notin \{w, w'\}$.

Abusing notation, let $\bar{\alpha}_\epsilon(\alpha)$ denote the aggregate action under mixture distribution $(1 - \epsilon)F + \epsilon\delta(w')$. Note that $\bar{\alpha}_\epsilon(\alpha) = \bar{\alpha}_\epsilon(\alpha')$ and $\bar{\alpha}_\epsilon(\alpha'') = \bar{\alpha}_\epsilon(\alpha''') = \bar{\alpha}_\epsilon(\alpha) + \epsilon(a''' - a'')$, where

$$\bar{\alpha}_\epsilon(\alpha) = (1 - \epsilon) \int_{\underline{w}}^{\bar{w}} \alpha(w) dF(w) + \epsilon a'' = (1 - \epsilon) \bar{\alpha}(\alpha) + \epsilon a''.$$

Given the mixture distribution $F^{\epsilon\epsilon} = (1 - \epsilon - \epsilon)F + \epsilon\delta(w) + \epsilon\delta(w')$ the definition of potential (3) implies that

$$\begin{aligned} & \lim_{\epsilon \rightarrow 0} \frac{(V(\alpha''', F^{\epsilon\epsilon}, \theta) - V(\alpha'', F^{\epsilon\epsilon}, \theta)) - (V(\alpha', F^{\epsilon\epsilon}, \theta) - V(\alpha, F^{\epsilon\epsilon}, \theta))}{\epsilon} \\ &= \Delta U(a', a, \bar{\alpha}_\epsilon(\alpha''), \theta, w) - \Delta U(a', a, \bar{\alpha}_\epsilon(\alpha), \theta, w) \\ &= \Delta U(a', a, \bar{\alpha}_\epsilon(\alpha) + \epsilon(a''' - a''), \theta, w) - \Delta U(a', a, \bar{\alpha}_\epsilon(\alpha), \theta, w) \end{aligned}$$

Since U is Lipschitz continuous, we can further divide this difference by ϵ and obtain

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \lim_{\varepsilon \rightarrow 0} \frac{(V(\alpha''', F^{\epsilon\varepsilon}, \theta) - V(\alpha'', F^{\varepsilon\varepsilon}, \theta)) - (V(\alpha', F^{\varepsilon\varepsilon}, \theta) - V(\alpha, F^{\varepsilon\varepsilon}, \theta))}{\varepsilon} \\ = (a''' - a'') \frac{\partial \Delta U(a', a, \bar{\alpha}(\alpha), \theta, w)}{\partial \bar{a}}. \end{aligned}$$

A similar argument shows that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \lim_{\epsilon \rightarrow 0} \frac{(V(\alpha''', F^{\epsilon\varepsilon}, \theta) - V(\alpha', F^{\epsilon\varepsilon}, \theta)) - (V(\alpha'', F^{\varepsilon\varepsilon}, \theta) - V(\alpha, F^{\varepsilon\varepsilon}, \theta))}{\epsilon} \\ = (a' - a) \frac{\partial \Delta U(a''', a'', \bar{\alpha}(\alpha), \theta, w')}{\partial \bar{a}}. \end{aligned}$$

Since the mixture distribution $F^{\epsilon\varepsilon}$ converges uniformly to F as $\varepsilon, \epsilon \rightarrow 0$ regardless of the order of limits, by the continuity of V , both limits must coincide, yielding the following *externality symmetry* condition: for all a, a', a'' and a''' , all α and all w, w' ,

$$(a''' - a'') \frac{\partial \Delta U(a', a, \bar{\alpha}(\alpha), \theta, w)}{\partial \bar{a}} = (a' - a) \frac{\partial \Delta U(a''', a'', \bar{\alpha}(\alpha), \theta, w')}{\partial \bar{a}}. \quad (26)$$

We next prove that (26) requires payoffs to be additively separable in \bar{a} and w . That is, they must take on the form $U(a, \bar{a}, \theta, w) = u(a, \bar{a}, \theta) + v(a, \theta, w) + u_0(\bar{a}, \theta, w)$. If we set $a'' = a$ and $a''' = a'$, (26) implies that

$$\frac{\partial \Delta U(a', a, \bar{\alpha}(\alpha), \theta, w)}{\partial \bar{a}} = \frac{\partial \Delta U(a', a, \bar{\alpha}(\alpha), \theta, w')}{\partial \bar{a}},$$

for all α and all w, w' . Hence, payoffs must be separable since the partial derivative of payoff differences w.r.t. the aggregate action is independent of types.

Consider next the linearity of $u(a, \bar{a}, \theta)$ w.r.t. a . First, if there are just two actions $a < a'$ in A then we can always write separable payoffs in a linear form $au(\bar{a}, \theta)$ by defining $u(\bar{a}, \theta) = \frac{1}{a' - a} (u(a', \bar{a}, \theta) - u(a, \bar{a}, \theta))$ and adding to u_0 the term $\frac{a'}{a' - a} u(a, \bar{a}, \theta) + \frac{a}{a' - a} u(a', \bar{a}, \theta)$. In general, for $w = w'$, (26) implies that

$$\frac{\partial}{\partial \bar{a}} \frac{\Delta U(a, a', \bar{a}, \theta, w)}{(a - a')} = \frac{\partial}{\partial \bar{a}} \frac{\Delta U(a''', a'', \bar{a}, \theta, w)}{(a''' - a'')},$$

i.e., $\frac{\partial}{\partial \bar{a}} \frac{\Delta U(a, a', \bar{a}, \theta, w)}{\Delta a}$ is independent of a and a' . Accordingly, we can write u as a linear function of a . Similarly, if A contains an interval, given the Lipschitz continuity of U , the cross-partial derivative $\frac{\partial^2 U(a, \bar{a}, \theta, w)}{\partial a \partial \bar{a}}$ is constant in a for almost all a in the interval and all \bar{a} , i.e., $u(a, \bar{a}, \theta)$ is linear in a .

Finally, since the existence of potential implies the above separability and linear-

ity restrictions, for Γ to be a weighted potential game, there must exist a function $\psi(\theta, w)$ such that $\psi(\theta, w)U(a, \bar{a}, \theta, w) = u(a, \bar{a}, \theta) + v(a, \theta, w) + u_0(\bar{a}, \theta, w)$. But this implies that $U(a, \bar{a}, \theta, w)$ satisfies (1), where $c(\theta, w) = 1/\psi(\theta, w) > 1/\zeta = \xi > 0$. That is, payoffs must be quasilinear.

“Only If” part: Given quasilinear payoffs consider the functional defined by (4). The change in V under mixture distribution F_w^ε when type w switches from action $\alpha(w)$ to a is given by

$$\int_{(1-\varepsilon)\bar{\alpha}(\alpha)+\varepsilon\alpha(w)}^{(1-\varepsilon)\bar{\alpha}(\alpha)+\varepsilon a} u(z, \theta) dz + \varepsilon (v(a, \theta, w) - v(\alpha(w), \theta, w)).$$

Dividing by ε and taking the limit, we obtain condition (3):

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \left(\int_{(1-\varepsilon)\bar{\alpha}(\alpha)+\varepsilon\alpha(w)}^{(1-\varepsilon)\bar{\alpha}(\alpha)+\varepsilon a} u(z, \theta) dz + \varepsilon (v(a, \theta, w) - v(\alpha(w), \theta, w)) \right) \\ &= \frac{\partial}{\partial \varepsilon} \int_{(1-\varepsilon)\bar{\alpha}(\alpha)+\varepsilon\alpha(w)}^{(1-\varepsilon)\bar{\alpha}(\alpha)+\varepsilon a} u(z, \theta) dz + v(a', \theta, w) - v(\alpha(w), \theta, w) \\ &= (a - \alpha(w))u(\bar{\alpha}(\alpha), \theta) + v(a', \theta, w) - v(\alpha(w), \theta, w) = \Delta U(a, \alpha(w), \bar{\alpha}(\alpha), \theta, w). \end{aligned}$$

That is, the change in V in the limit coincides with the change in payoffs for type w , up to scaling by a function $c(\theta, w)$ so V is a (weighted) potential of Γ . \square

B.2 Proofs of Results in Subsection 4.1

Proof of Proposition 2. To prove the first part we show that the problem of maximizing potential coincides with the problem of maximizing ex-ante payoffs under marginal beliefs given by (5). Maximizing potential implies finding a strategy profile that solves

$$\max_{\alpha \in \mathcal{A}} \int_0^{\bar{\alpha}(\alpha)} u(z, \theta) dz + \int_{\underline{w}}^{\bar{w}} v(\alpha(w), \theta, w) dF(w). \quad (27)$$

Consider the change of variable $z = \bar{\alpha}(\alpha, [w, \bar{w}]) (1 - F(w))$ to the first integral. Differentiating $\bar{\alpha}(\alpha, [w, \bar{w}])$ w.r.t. w we obtain $dz = -\alpha(w) f(w) dw$. In addition, $0 = \bar{\alpha}(\alpha, [\bar{w}, \bar{w}])$ and $\bar{\alpha}(\alpha) = \bar{\alpha}(\alpha, [\underline{w}, \bar{w}])$ which leads to objective function

$$\int_{\underline{w}}^{\bar{w}} \alpha(w) u(\bar{\alpha}(\alpha, [w, \bar{w}]) (1 - F(w)), \theta) f(w) dw + \int_{\underline{w}}^{\bar{w}} v(\alpha(w), \theta, w) dF(w).$$

To prove that α_P must be a NE assume, by way of contradiction, that α_P maximizes potential but that there is a closed set of types W of positive measure such

that, for each $w \in W$, there exists some $\alpha(w)$ such that $\Delta U(\alpha_P(w), \alpha(w), \bar{\alpha}(\alpha), \theta, w) < 0$, implying that α_P is not a NE. Let $\alpha(w') = \alpha_P(w')$ for all $w' \notin W$. The difference in potential between α_P and α can be written as

$$V(\alpha_P, \theta) - V(\alpha, \theta) = \int_0^{\bar{\alpha}(\alpha_P) - \bar{\alpha}(\alpha)} u(\bar{\alpha}(\alpha) + z, \theta) dz + \int_W (v(\alpha_P(w), \theta, w) - v(\alpha(w), \theta, w)) dF(w).$$

Let $z(w) = \int_{w' \in W} \mathbf{1}_{\{w' \geq w\}} (\alpha_P(w') - \alpha(w')) dF(w')$. Differentiating w.r.t. w we obtain $dz = -(\alpha_P(w) - \alpha(w)) dw$. Also, $z(\min W) = \bar{\alpha}(\alpha_P) - \bar{\alpha}(\alpha)$ and $z(\max W) = 0$. Hence, applying a change of variable we get that

$$\begin{aligned} V(\alpha_P, \theta) - V(\alpha, \theta) &= \int_W (\alpha_P(w) - \alpha(w)) u\left(\bar{\alpha}(\alpha) + \int_W \mathbf{1}_{\{w' \geq w\}} (\alpha_P(w') - \alpha(w')) dF(w'), \theta\right) dw \\ &\quad + \int_W (v(\alpha_P(w), \theta, w) - v(\alpha(w), \theta, w)) dF(w) \\ &= \int_W \Delta U\left(\alpha_P(w), \alpha(w), \bar{\alpha}(\alpha) + \int_W \mathbf{1}_{\{w' \geq w\}} (\alpha_P(w') - \alpha(w')) dF(w'), \theta, w\right) dw. \end{aligned}$$

Note that, as the mass of W vanishes, $\int_W \mathbf{1}_{\{w' \geq w\}} (\alpha_P(w') - \alpha(w')) dF(w')$ goes to zero and $\bar{\alpha}(\alpha)$ uniformly converges to $\bar{\alpha}(\alpha_P)$, implying that the integrand in the above expression converges to $\Delta U(\alpha_P(w), \alpha(w), \bar{\alpha}(\alpha_P), \theta, w)$ for all $w \in W$. Since ΔU is Lipschitz continuous and $\Delta U(\alpha_P(w), \alpha(w), \bar{\alpha}(\alpha_P), \theta, w) < 0$ for all $w \in W$, we can always find a set of W with small enough probability mass so that

$$\Delta U\left(\alpha_P(w), \alpha(w), \bar{\alpha}(\alpha) + \int_W \mathbf{1}_{\{w' \geq w\}} (\alpha_P(w') - \alpha(w')) dF(w'), \theta, w\right) < 0$$

for all $w \in W$. But this implies that $V(\alpha_P, \theta) - V(\alpha, \theta) < 0$, a contradiction.

Finally, we argue that set of solutions to the problem of maximizing potential given by (27) is non-empty. To do so we first show that, given any finite partition of the space of types into disjoint intervals, the set of potential maximizers is non-empty if we restrict attention to strategy profiles that map types in each of those intervals to the same action. Subsequently, we use the fact that measurable functions can be approximated by simple functions to show the existence of a measurable strategy profile that maximizes potential.

Let $\{W_j\}_{j=1}^n$ be a partition of $[\underline{w}, \bar{w}]$ into intervals W_j of the same length and define the set of strategy profiles that assign the same action to types in each W_j as

$$\mathcal{A}^n = \{\alpha : \alpha(w) = a_j \in A \text{ for all } w \in W_j, j = 1, \dots, n\}.$$

Since each W_j is measurable \mathcal{A}^n is a subset of \mathcal{A} . Hence, V is well-defined and continuous in \mathcal{A}^n . In addition, \mathcal{A}^n it coincides with the space A^n , which is compact in the product topology since A is compact. Hence, by Weiertrass extreme value theorem, $\max_{\alpha \in \mathcal{A}^n} V(\alpha, \theta)$ has a solution for any $n < \infty$. Let $V^n := \max_{\alpha \in \mathcal{A}^n} V(\alpha, \theta)$.

Next, note that the sequence $\{V^{2n}\}$ converges as $n \rightarrow \infty$. This is because the subsets in $\{W_j\}_{j=1}^n$ are unions of subsets in $\{W_j\}_{j=1}^{2n}$, implying that $\mathcal{A}^n \subset \mathcal{A}^{2n}$ and thus $V^{2n} \geq V^n$ for all n . Hence, $\{V^{2n}\}$ is a monotone bounded sequence so it must converge. Accordingly, by Lipschitz continuity of u and v and the compactness of \mathcal{A}^{2n} , convergence of $\{V^{2n}\}$ implies that we can find a convergent sequence $\{\alpha^{2n}\}$ with $\alpha^{2n} \in \mathcal{A}^{2n}$ for all n such that $\lim V^{2n} = \lim V(\alpha^{2n}, \theta)$. But since the sets \mathcal{A}^n are subsets of simple functions in \mathcal{A} (i.e., they take on finitely many values), $\{\alpha^{2n}\}$ must converge to a measurable function, i.e. to a strategy in \mathcal{A} . This shows that there exists $\alpha \in \mathcal{A}$ such that $\lim V^n \leq V(\alpha, \theta)$.

To finish the proof we argue that $\lim V^n = \max_{\alpha \in \mathcal{A}} V(\alpha, \theta)$. Assume otherwise that there is α' such that $V(\alpha', \theta) > \lim V^n$. Since every measurable function can be approximated by a sequence of simple functions, by the continuity of V we can find a high enough n' such that $V(\alpha', \theta)$ is arbitrarily close to $V(\alpha^{n'}, \theta)$ for some $\alpha^{n'} \in \mathcal{A}^{n'}$. But then we have that $V^{n'} \geq V(\alpha^{n'}, \theta) > \lim V^n \geq V^{n'}$, a contradiction. \square

Proof of Proposition 3. Existence of at least one strategy profile α_P that maximizes potential is guaranteed by Proposition 2. In addition, NE strategy profiles must be (weakly) increasing in w since payoffs exhibit strictly increasing differences in a and w . Hence, any α_P satisfying (5) must be monotone given that it is also a NE.

Next, we show that α_P is unique except perhaps in a countable subset of Θ . We do so by showing that, if there is more than one potential maximizer at some θ , then there exist $\theta' < \theta$ and $\theta'' > \theta$ such that there is only one potential maximizer in $(\theta', \theta) \cup (\theta, \theta'')$. Since Θ can only be partitioned in a countable number of non-degenerate intervals then the set of θ at which there are multiple potential maximizers must be countable, i.e., must have Lebesgue measure zero.

First, note that if there exist two maximizers α_1 and α_2 then the monotonicity of NE implies that $\alpha_2(w) \geq \alpha_1(w)$ for all w , with strict inequality for a positive mass of types. In turn, this implies that $\bar{\alpha}(\alpha_2, [w, \bar{w}]) \geq \bar{\alpha}(\alpha_1, [w, \bar{w}])$ for all w , with strict inequality for a positive mass of types.

Second, note that if α_1 and α_2 maximize potential for a given θ , they satisfy (5) and thus we must have that

$$\int_w^{\bar{w}} \left(U(\alpha_2(w), \bar{\alpha}(\alpha_2, [w, \bar{w}])(1 - F(w)), \theta, w) - U(\alpha_1(w), \bar{\alpha}(\alpha_1, [w, \bar{w}])(1 - F(w)), \theta, w) \right) dF(w) = 0.$$

By Assumption 1 the integrand is strictly increasing in θ for all w s.t. $\alpha_2(w) > \alpha_1(w)$

or $\bar{\alpha}(\alpha_2, [w, \bar{w}]) > \bar{\alpha}(\alpha_1, [w, \bar{w}])$. Hence, an infinitesimal increase in θ would make the LHS go up, leading, by the Lipschitz continuity of U , to a unique profile maximizing potential $\alpha'_2(w) \geq \alpha_2(w)$ for all w . Similarly, an infinitesimal drop in θ would lead to a unique potential-maximizing profile $\alpha'_1(w) \leq \alpha_1(w)$ for all w . These arguments also apply to the case of more than two profiles maximizing potential at θ , i.e., an infinitesimal increase (decrease) $d\theta$ leads to a unique potential-maximizing profile that is weakly higher (lower) than the highest (lowest) potential maximizer at θ . Accordingly, there is a unique maximizer in an open neighborhood $(\theta', \theta) \cup (\theta, \theta'')$. \square

B.3 Proofs of Results in Subsection 4.2

The proofs of [Propositions 4](#) and [5](#) first focus on the uniform prior case and then resort to the following lemma about the uniform convergence of individual beliefs as noise vanishes to extend the results to any well-defined prior.

Lemma 6. *Let $J_{w'|w}(s'|s; \nu, \phi)$ denote the cdf of signals of type w' agents conditional on an agent of type w receiving signal s , for given noise level ν and common prior ϕ . Given any $s \in [\inf \Theta + \nu, \sup \Theta - \nu]$ and any sequence s^ν such that $\frac{s-s^\nu}{\nu} = c$ for some constant $|c| < 1$, as $\nu \rightarrow 0$, $|J_{w'|w}(s^\nu|s; \nu, \phi) - J_{w'|w}(s^\nu|s; \nu, U[\inf \Theta, \sup \Theta])|$ converges uniformly to zero.*

Proof of Lemma 6. Let s_w denote the random variable representing the signals received by agents of type w . The beliefs about s_w of an agent of type w conditional on receiving s can be expressed as

$$J_{w'|w}(s'|s; \nu, \phi) = \frac{\Pr(s'_w < s', s_w = s)}{\Pr(s_w = s)} = \frac{\int_{s-\nu/2}^{s+\nu/2} H_{w'}\left(\frac{s'-\theta}{\nu}\right) h_w\left(\frac{s-\theta}{\nu}\right) \frac{1}{\nu} \phi(\theta) d\theta}{\int_{s-\nu/2}^{s+\nu/2} h_w\left(\frac{s-\theta}{\nu}\right) \frac{1}{\nu} \phi(\theta) d\theta}$$

Using the change of variable $\theta' = \frac{s-\theta}{\nu}$, for any $s' = s - c\nu$ we have that

$$\begin{aligned} J_{w'|w}(s'|s; \nu, \phi) &= \frac{\int_{-1/2}^{1/2} H_{w'}(\theta' - c) h_w(\theta') \phi(s - \nu\theta') d\theta'}{\int_{-1/2}^{1/2} h_w(\theta') \phi(s - \nu\theta') d\theta'} \\ &\rightarrow \int_{-1/2}^{1/2} H_{w'}(\theta' - c) h_w(\theta') d\theta' = J_{w'|w}(s'|s; \nu, U[\inf \Theta, \sup \Theta]), \end{aligned}$$

as $\nu \rightarrow 0$. Since $J_{w'|w}$ is a cdf and the limit is continuous, pointwise convergence implies uniform convergence. \square

Proof of Proposition 4. The proof logic is as follows. First, we argue for the uniform prior case that, given any $\nu > 0$, the set of equilibrium strategy profiles has a largest and a smallest element, each involving monotone strategies. Second, we show that there is at most one equilibrium in monotone strategies, up to differences in behavior at cutoff signals, so the smallest and largest equilibria are essentially the same. These arguments extend to the non-uniform prior case by the uniform convergence of beliefs (Lemma 6).

Consider the game in which we fix the profile of signal realizations and agents choose actions in A after observing their own signals. Given Assumption 1 the game satisfies the conditions of Theorem 5 in Milgrom and Roberts (1990).²⁰ Accordingly, it has a smallest equilibrium α^l and a largest equilibrium α^m such that any equilibrium profile α satisfies $\alpha^l(w, s) \leq \alpha(w, s) \leq \alpha^m(w, s)$.

In addition, fixing the actions of all agents, an agent's difference in expected payoffs conditional on s from choosing a versus $a' < a$ is increasing in s since the aggregate action is kept fixed while θ is higher (in expectation) at higher signal profiles. That is, expected payoffs exhibit increasing differences w.r.t. a and the profile of signals for every w . Hence, Theorem 6 in Milgrom and Roberts (1990) applies: the smallest and largest equilibria are nondecreasing w.r.t. the profile of signals. Since an agent's strategy can only depend on her own signal, this implies that α^l and α^m are monotone functions of s . A similar argument applies to monotonicity w.r.t. w .

To show that there is at most one equilibrium in monotone strategies, we first establish the following translation result. Given $\delta > 0$, let α_δ represent a "rightward shift" of strategy profile α defined by

$$\alpha_\delta(w, s) = \begin{cases} \alpha(w, \inf \Theta - \nu/2) & s < \inf \Theta - \nu/2 + \delta \\ \alpha(w, s - \delta) & s \geq \inf \Theta - \nu/2 + \delta. \end{cases}$$

The next lemma shows that if we simultaneously switch agents' strategies from α to α_δ and their signals from s to $s + \delta$ then an agent's conditional expectation of payoff differences between a higher action a and a lower action $a' < a$ strictly increases. [We omit the dependence of α on ν to ease notation.]

Lemma 7. *There exists $\bar{\nu} > 0$ such that, for any $\alpha \in \mathcal{A}$, if $\nu < \bar{\nu}$ then*

$$E[\Delta U(a, a', \bar{\alpha}(\alpha_\delta), \theta, w)|s + \delta] - E[\Delta U(a, a', \bar{\alpha}(\alpha), \theta, w)|s] \geq K(a - a')\delta \quad (28)$$

for all actions a and $a' < a$, all $w \in [\underline{w}, \bar{w}]$, all $s \in [\underline{\theta} - \nu/2, \bar{\theta} + \nu/2]$ and all $\delta \in (0, \sup \Theta - s - \nu/2]$.

²⁰The theorem applies given that the set \mathcal{A} of measurable functions from $[\underline{w}, \bar{w}]$ to A is a lattice.

Proof. For all $\nu < \bar{\nu} := \min\{\underline{\theta} - \inf \Theta, \sup \Theta - \bar{\theta}\}$, the support of the distributions of θ and other player signals conditional on $s \in [\underline{\theta} - \nu/2, \bar{\theta} + \nu/2]$ are respectively $[s - \nu/2, s + \nu/2]$ and $[s - \nu, s + \nu]$. In such a case, the conditional density of θ is $h_w\left(\frac{s-\theta}{\nu}\right)$. Also notice that, conditional on θ , the signals of other agents are independent of s , with densities $h_{w'}\left(\frac{s'-\theta}{\nu}\right)$. Applying the exact LLN within type, the aggregate action given θ is given by

$$\bar{\alpha}(\alpha) = \int_{\underline{w}}^{\bar{w}} \int_{\theta-\nu/2}^{\theta+\nu/2} \alpha(w', s') h_{w'}\left(\frac{s'-\theta}{\nu}\right) ds' dF(w'). \quad (29)$$

By [Assumption 1](#), for any $s \in [\underline{\theta} - \nu/2, \bar{\theta} + \nu/2]$ and $\delta \in (0, \sup \Theta - s - \nu/2]$ we obtain the following inequality using the changes of variables $\theta' = \theta + \delta$ and $s'' = s' + \delta$:

$$\begin{aligned} & E[\Delta U(a, a', \bar{\alpha}(\alpha), \theta, w) | s] + K(a - a')\delta = \\ & \int_{s-\nu/2}^{s+\nu/2} \Delta U \left(a, a', \int_{\underline{w}}^{\bar{w}} \int_{\theta-\nu/2}^{\theta+\nu/2} \alpha(w', s') h_{w'}\left(\frac{s'-\theta}{\nu}\right) ds' dF(w'), \theta, w \right) h_w\left(\frac{s-\theta}{\nu}\right) d\theta + K(a - a')\delta \\ & \leq \int_{s-\nu/2}^{s+\nu/2} \Delta U \left(a, a', \int_{\underline{w}}^{\bar{w}} \int_{\theta-\nu/2}^{\theta+\nu/2} \alpha(w', s') h_{w'}\left(\frac{s'-\theta}{\nu}\right) ds' dF(w'), \theta + \delta, w \right) h_w\left(\frac{s-\theta}{\nu}\right) d\theta \\ & = \int_{s+\delta-\nu/2}^{s+\delta+\nu/2} \Delta U \left(a, a', \int_{\underline{w}}^{\bar{w}} \int_{\theta'-\delta-\nu/2}^{\theta'-\delta+\nu/2} \alpha(w', s') h_{w'}\left(\frac{s'+\delta-\theta'}{\nu}\right) ds' dF(w'), \theta', w \right) h_w\left(\frac{s+\delta-\theta'}{\nu}\right) d\theta' \\ & = \int_{s+\delta-\nu/2}^{s+\delta+\nu/2} \Delta U \left(a, a', \int_{\underline{w}}^{\bar{w}} \int_{\theta'+\nu/2}^{\theta'+\nu/2} \alpha(w', s'' - \delta) h_{w'}\left(\frac{s''-\theta'}{\nu}\right) ds'' dF(w'), \theta', w \right) h_w\left(\frac{s+\delta-\theta'}{\nu}\right) d\theta' \\ & = E[\Delta U(a, a', \bar{\alpha}(\alpha_\delta), \theta, w) | s + \delta].^{21} \quad \square \end{aligned}$$

Next, note that if α is an equilibrium then $\alpha(w, s) = 0$ for all $s \leq \underline{\theta} - \nu/2$ and $\alpha(w, s) = a_{max}$ if $s > \bar{\theta} + \nu/2$. This is because, for any action $a > 0$, [Assumption 2](#)

²¹The change of variable $\theta' = \theta + \delta$ works as long as the upper integration limit is well-defined, i.e., $s + \delta + \nu/2 \leq \sup \Theta$, which is guaranteed by $\delta \leq \sup \Theta - s - \nu/2$. The change of variable $s'' = s' + \theta$ works as long as $\theta' - \delta \geq \inf \Theta$, otherwise $\alpha(s'' - \delta, w')$ is not well-defined. Since $\theta' \geq s + \delta - \nu/2$ and $s \geq \underline{\theta} - \nu/2$ we have that $\theta' - \delta \geq \underline{\theta} - \nu$. Hence, $\theta' - \delta \geq \inf \Theta$ for any $\nu < \bar{\nu}$.

implies that $\Delta U(a, 0, \theta, w) < 0$ for all $\theta < \underline{\theta}$. Hence, since $\theta \leq s + \nu/2$, it must be that $E[\Delta U(a, 0, \bar{\alpha}(\alpha), \theta, w)|s] < 0$ for all $s < \underline{\theta} - \nu/2$. A symmetric argument applies to signals above $\bar{\theta} + \nu/2$.

We finish the proof by arguing that $\alpha^l(w, s) = \alpha^m(w, s)$ for almost all (w, s) using the above translation result. Assume first, by way of contradiction, that $\alpha^l(w, s) < \alpha^m(w, s)$ for some signal s and some type w and that there exists a signal shift $\delta > 0$ such that $\alpha^l(w, s + \delta) = \alpha^m(w, s)$ or, if $\alpha^m(w, \cdot)$ is discontinuous at s , $\alpha^l(w, s + \delta) \in [\liminf \alpha^m(w, s), \limsup \alpha^m(w, s)]$. Note that the monotonicity of $\alpha^l(w, \cdot)$ and $\alpha^m(w, \cdot)$ means that their lim inf and lim sup exist. Next, consider among all pairs (w, s) at which the two equilibria differ the largest signal shift $\hat{\delta}$ that would be required to make $\alpha^l(w, s)$ ‘equal’ to $\alpha^m(w, s)$. That is $\hat{\delta}$ is given by

$$\hat{\delta} = \max \left\{ \delta : \alpha^l(w, s + \delta) \in [\liminf \alpha^m(w, s), \limsup \alpha^m(w, s)] \right. \\ \left. \text{for some } (w, s) \text{ s.t. } \alpha^l(w, s + \delta) \neq \alpha^m(w, s + \delta) \right\}.$$

It is straightforward to check that $\alpha^m(w, s - \hat{\delta}) \leq \alpha^l(w, s)$ for all s, w . Let (\hat{w}, \hat{s}) be one of the signal-type pairs associated with $\hat{\delta}$. Note that, by the above argument, we must have that $\hat{s} + \hat{\delta} \in [\underline{\theta} - \nu/2, \bar{\theta} + \nu/2]$, otherwise $\alpha^l(\hat{s} + \hat{\delta}, \hat{w}) = \alpha^m(\hat{s} + \hat{\delta}, \hat{w}) \in \{0, a_{max}\}$. Also note that if α is an equilibrium we must have that $E[\Delta U(a, \alpha(w, s), \bar{\alpha}(\alpha), \theta, w)|s] \leq 0$ for all $a \in A$. Hence, by [Lemma 7](#), we arrive to the following contradiction:

$$0 \leq E[\Delta U(\alpha^m(\hat{w}, \hat{s}), \alpha^l(\hat{w}, \hat{s}), \bar{\alpha}(\alpha^m), \theta, \hat{w})|\hat{s}] < E[\Delta U(\alpha^m(\hat{w}, \hat{s}), \alpha^l(\hat{w}, \hat{s}), \bar{\alpha}(\alpha_{\hat{s}}^m), \theta, \hat{w})|\hat{s} + \hat{\delta}] \\ \leq E[\Delta U(\alpha^m(\hat{w}, \hat{s}), \alpha^l(\hat{w}, \hat{s}), \bar{\alpha}(\alpha^l), \theta, \hat{w})|\hat{s} + \hat{\delta}] \leq E[\Delta U(\alpha^m(\hat{w}, \hat{s}), \alpha^l(\hat{w}, \hat{s} + \hat{\delta}), \bar{\alpha}(\alpha^l), \theta, \hat{w})|\hat{s} + \hat{\delta}] \\ \leq 0.$$

The third inequality comes from [Assumption 1](#) and the fact that $\alpha^m(\hat{w}, \hat{s}) > \alpha^l(\hat{w}, \hat{s})$ and $\bar{\alpha}(\alpha^l) \geq \bar{\alpha}(\alpha_{\hat{s}}^m)$ since $\alpha^m(w, s - \hat{\delta}) \leq \alpha^l(w, s)$ for all s, w . The last two inequalities follow from $\alpha^l(\hat{w}, \hat{s} + \delta)$ being a best response of type \hat{w} to α^l under $\hat{s} + \hat{\delta}$.

Hence, the only possibility left for $\alpha^l(w, s) < \alpha^m(w, s)$ is that there is no signal shift $\delta > 0$ that satisfies $\alpha^l(w, s + \delta) \in [\liminf \alpha^m(w, s), \limsup \alpha^m(w, s)]$. Given that both $\alpha^l(w, \cdot)$ and $\alpha^m(w, \cdot)$ are increasing this can only happen in the set of s at which they are discontinuous, which has zero measure. \square

Proof of [Proposition 5](#). We first prove under a uniform prior that equilibrium strategy profiles essentially converge to some strategy profile α_G . Specifically, we show that if α^ν is any sequence of equilibrium strategy profiles in the global game indexed by $\nu \rightarrow 0$, given any $\varepsilon > 0$ then there exists $\bar{\nu} > 0$ such that, for all $\bar{\nu} > \nu > \nu' > 0$, $|\alpha^\nu(w, s) - \alpha^{\nu'}(w, s)| < \varepsilon$ for almost all s and all w .

Abusing notation, let $\bar{\alpha}^\nu(\alpha; \theta)$ denote the aggregate action when players follow strategy profile α , the noise level is ν and the true parameter is θ .

For any monotone strategy profile α let $S_d(\alpha)$ be the set of signals at which $\alpha(w, \cdot)$ is discontinuous for a positive measure of types w . Since $\alpha(w, \cdot)$ is monotone, $S_d(\alpha)$ is at most countable, i.e., it has zero measure for all $\nu > 0$. Recall that, conditional on an agent receiving s , the profile of realized signals must lie inside $[s - \nu, s + \nu]$. Hence, given that signals and types are continuously distributed, the mass of agents with signals that belong to $S_d(\alpha)$ conditional on a player receiving signal $s \notin S_d(\alpha)$ must be zero for all $\nu \geq 0$, that is, even in the limit since $[s - \nu, s + \nu] \rightarrow \{s\}$ as $\nu \rightarrow 0$. But this implies that the aggregate action $\bar{\alpha}^\nu(\alpha; \theta)$ converges uniformly to

$$\bar{\alpha}(\alpha; s) := \int_{\underline{w}}^{\bar{w}} \alpha(w, s) dF(w).$$

To see why, notice that the aggregate action given for any given θ is

$$\bar{\alpha}^\nu(\alpha; \theta) = \int_{\underline{w}}^{\bar{w}} \int_{-1/2}^{1/2} \alpha(w, \theta + \nu\eta) h_w(\eta) d\eta dF(w).$$

Since $\theta \in [s - \nu/2, s + \nu/2]$, by the monotonicity of α we can bound $\bar{\alpha}^\nu(\alpha; \theta)$:

$$\int_{\underline{w}}^{\bar{w}} \int_{-1/2}^{1/2} \alpha(w, s - \nu/2 + \nu\eta) h_w(\eta) d\eta dF(w) \leq \bar{\alpha}^\nu(\alpha; \theta) \leq \int_{\underline{w}}^{\bar{w}} \int_{-1/2}^{1/2} \alpha(w, s + \nu/2 + \nu\eta) h_w(\eta) d\eta dF(w).$$

Since $s \pm \nu/2 + \nu\eta \in [s - \nu, s + \nu]$ for all η and $s \notin S_d(\alpha)$ the set of pairs $(w, s \pm \nu/2 + \nu\eta)$ at which α is discontinuous has zero measure so both bounds must converge to $\bar{\alpha}(\alpha; s)$. Moreover, $\alpha(w, s - \nu/2 + \nu\eta)$ and $\alpha(w, s + \nu/2 + \nu\eta)$ respectively increase and decrease as $\nu \rightarrow 0$ for all $\eta \in (-1/2, 1/2)$, implying that the lower bound monotonically increases and the upper bound monotonically decreases. That is, the convergence of $\bar{\alpha}^\nu(\alpha; \theta)$ to $\bar{\alpha}(\alpha, s)$ must be uniform.

In turn, by the Lipschitz continuity of ΔU , the uniform convergence of $\bar{\alpha}^\nu(\alpha; \theta)$ implies that expected payoff differences between any two actions conditional on receiving $s \notin S_d(\alpha)$ uniformly converge as $\nu \rightarrow 0$ for any fixed monotone profile α . That is, for all $s \notin S_d(\alpha)$ and all w and all $\varepsilon > 0$, there is $\hat{\nu} > 0$ such that

$$|E_\nu(\Delta U(\alpha, a', \bar{\alpha}^\nu(\alpha; \theta), \theta, w)|s) - E_{\nu'}(\Delta U(\alpha, a', \bar{\alpha}^{\nu'}(\alpha, \theta), \theta, w)|s)| < \varepsilon$$

for all $\nu, \nu' < \hat{\nu}$, where E_ν denotes the expectation operator under noise level ν .

Equipped with this result we next argue that the equilibrium profiles must con-

verge. We do so by showing that for all $\nu, \nu' < \hat{\nu}$ the largest signal shift needed to make α^ν equal to $\alpha^{\nu'}$ is $O(\varepsilon)$. To account for discontinuities in $\alpha^{\nu'}$, define such a signal shift as

$$\hat{\delta} = \max \left\{ \delta : \alpha^\nu(w, s + \delta) \in [\liminf \alpha^{\nu'}(s, w), \limsup \alpha^{\nu'}(s, w)] \right. \\ \left. \text{for some } (w, s) \text{ s.t. } \alpha^\nu(w, s + \delta) \neq \alpha^{\nu'}(w, s + \delta) \right\}.$$

Assume that $\hat{\delta} > 0$ (otherwise we can apply a similar argument by switching α^ν and $\alpha^{\nu'}$). Note that $\alpha^{\nu'}(w, s - \hat{\delta}) \leq \alpha^\nu(w, s)$ for all s, w . Fix signal-type pair (\hat{w}, \hat{s}) associated with the signal shift $\hat{\delta}$, implying that $\alpha^{\nu'}(w, s) > \alpha^\nu(w, s)$. By [Lemma 7](#) and the fact that $\alpha^\nu, \alpha^{\nu'}$ are respectively the equilibrium strategies under noise levels ν, ν' , we have that

$$\begin{aligned} 0 &\leq E_{\nu'}[\Delta U(\alpha^{\nu'}(\hat{w}, \hat{s}), \alpha^\nu(\hat{w}, \hat{s}), \bar{a}(\alpha^{\nu'}), \theta, \hat{w}) | \hat{s}] \\ &\leq E_{\nu'}[\Delta U(\alpha^{\nu'}(\hat{w}, \hat{s}), \alpha^\nu(\hat{w}, \hat{s}), \bar{a}(\alpha^{\nu'}), \theta, \hat{w}) | \hat{s} + \hat{\delta}] - K(\alpha^{\nu'}(\hat{w}, \hat{s}) - \alpha^\nu(\hat{w}, \hat{s}))\hat{\delta} \\ &\leq E_{\nu'}[\Delta U(\alpha^{\nu'}(\hat{w}, \hat{s}), \alpha^\nu(\hat{w}, \hat{s}), \bar{a}(\alpha^\nu), \theta, \hat{w}) | \hat{s} + \hat{\delta}] - K(\alpha^{\nu'}(\hat{w}, \hat{s}) - \alpha^\nu(\hat{w}, \hat{s}))\hat{\delta} \\ &\leq E_{\nu'}[\Delta U(\alpha^{\nu'}(\hat{w}, \hat{s}), \alpha^\nu(\hat{w}, \hat{s}), \bar{a}(\alpha^\nu), \theta, \hat{w}) | \hat{s} + \hat{\delta}] + \varepsilon - K(\alpha^{\nu'}(\hat{w}, \hat{s}) - \alpha^\nu(\hat{w}, \hat{s}))\hat{\delta} \\ &\leq E_{\nu'}[\Delta U(\alpha^{\nu'}(\hat{w}, \hat{s}), \alpha^\nu(\hat{w}, \hat{s} + \hat{\delta}), \bar{a}(\alpha^\nu), \theta, \hat{w}) | \hat{s} + \hat{\delta}] + \varepsilon - K(\alpha^{\nu'}(\hat{w}, \hat{s}) - \alpha^\nu(\hat{w}, \hat{s}))\hat{\delta} \\ &\leq \varepsilon - K(\alpha^{\nu'}(\hat{w}, \hat{s}) - \alpha^\nu(\hat{w}, \hat{s}))\hat{\delta}. \end{aligned}$$

Since $\alpha^{\nu'}(\hat{w}, \hat{s}) - \alpha^\nu(\hat{w}, \hat{s})$ is bounded by a_{max} then $\hat{\delta} \leq \frac{\varepsilon}{Ka_{max}}$. That is, equilibrium strategies converge to some limit profile $\alpha_G(w, s)$ for almost all s and w .

Finally, notice that the convergence of equilibrium strategies and the uniform convergence of expected payoff differences, combined with the fact that $s \rightarrow \theta$ as $\nu \rightarrow 0$, imply that

$$\begin{aligned} \lim_{\nu \rightarrow 0} E_{\nu'}(\Delta U(\alpha^\nu(w, s), a, \bar{a}^\nu(\alpha^\nu; \theta), \theta, w) | s) &= E(\Delta U(\alpha_G(w, s), a, \bar{a}(\alpha_G(\cdot, s)), \theta, w) | s = \theta) \\ &= \Delta U(\alpha_G(w, \theta), a, \bar{a}(\bar{a}(\alpha_G(\cdot, \theta))), \theta, w), \end{aligned}$$

for almost all s . Since α^ν is an equilibrium on Γ^ν , we have that

$$E_{\nu'}(\Delta U(\alpha^\nu(w, s), a, \bar{a}^\nu(\alpha^\nu; \theta), \theta, w) | s) \geq 0,$$

for all s and all w . By the continuity of ΔU the above convergence implies that

$$\Delta U(\alpha_G(w, \theta), a, \bar{a}(\alpha_G(\cdot, \theta))), \theta, w) \geq 0,$$

for almost all θ and all w . That is, α_G is a NE of Γ for almost all θ . By [Lemma 6](#) the result extends to the non-uniform prior case. \square

B.4 Proofs of Results in [Subsection 4.3](#)

Proof of [Theorem 1](#). To prove the equivalence between α_P and α_G for almost all θ , we need to focus on the case when Γ has multiple NE but a unique potential maximizer. This is because (i) when Γ has a unique NE it must coincide with both α_P and α_G for almost all θ by [Propositions 3](#) and [5](#); and (ii) these propositions also imply that the set of θ at which Γ has multiple potential maximizers or at which the global games selection is not unique has measure zero.

Focus first on the uniform-prior case. The proof applies the Generalized Laplacian property ([Lemma 3](#)) to show that if α_P is the unique potential maximizer then

$$\lim_{\nu \rightarrow 0} E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}^\nu(\alpha_P; \theta), \theta, w) | s) > 0$$

for all $a \neq \alpha_P(w, \theta)$, all $s \in (\inf \Theta + \nu/2, \sup \Theta - \nu/2)$ and almost all w , where $\bar{\alpha}^\nu(\alpha_P; \theta)$ represents the aggregate action given θ when agents follow strategies $\alpha_P(w, s)$. But, as shown in the proof of [Proposition 5](#), these are the conditions that fully characterize the limit equilibrium in the global game, which is essentially unique by [Proposition 4](#). Hence, it must be that $\alpha_P = \alpha_G$ for almost all θ and almost all w .

Fix any $\theta \in (\inf \Theta, \sup \Theta)$ under which there is a unique potential maximizer. For any given any action $a \in A$ and any subset of types W with positive measure such that $\alpha_P(w, \theta) > a$ for all $w \in W$, let $\alpha(w) = a$ if $w \in W$ and $\alpha(w) = \alpha_P(w, \theta)$ if $w \notin W$. Since α_P is the unique potential maximizer, we have that

$$V(\alpha_P, \theta) - V(\alpha, \theta) = \int_{\bar{\alpha}(\alpha)}^{\bar{\alpha}(\alpha_P(\cdot, \theta))} u(z, \theta) dz + \int_W (v(\alpha_P(w, \theta), \theta, w) - v(a, \theta, w)) dF(w) > 0.$$

Let $\alpha(w, s)$ represent the monotone strategy in the global game in which agents of types in W switch from a to $\alpha_P(w, \theta)$ using cutoff function $\kappa(w) = \theta$ for all $w \in W$; while types in $W^C := [\underline{w}, \bar{w}] \setminus W$ choose $\alpha(w, s) = \alpha_P(w, \theta)$ for all s . We can express their expected payoff differences conditional *both* on $s = \kappa(w)$ and on θ as follows:

$$\begin{aligned}
& E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}^\nu(\alpha; \theta), \theta, w) | s = \theta, \theta) \\
&= \int_a^{\bar{\alpha}(\alpha_P(\cdot, \theta), W)} (\alpha_P(w, \theta) - a) u(\bar{\alpha}(\alpha_P(\cdot, \theta), W^C) F(W^C) + z F(W), \theta) dG_w(z | \kappa; \alpha, W) \\
&+ (v(\alpha_P(w, \theta), \theta, w) - v(a, \theta, w)),
\end{aligned}$$

where z represents the aggregate action of agents with types in W and $F(W^C)$, $F(W)$ respectively denote the probability mass of W^C and W . Integrating the above expression over types in W and applying [Lemma 3](#) we obtain

$$\begin{aligned}
& \int_W E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}^\nu(\alpha; \theta), \theta, w) | s = \theta, \theta) dF(w | w \in W) \\
&= \int_a^{\bar{\alpha}(\alpha_P(\cdot, \theta), W)} u(\bar{\alpha}(\alpha_P(\cdot, \theta), W^C) F(W^C) + z F(W), \theta) dz \\
&+ \int_W (v(\alpha_P(w, \theta), \theta, w) - v(a, \theta, w)) dF(w | w \in W). \tag{30}
\end{aligned}$$

Applying the change of variable $z' = \bar{\alpha}(\alpha_P(\cdot, \theta), W^C) F(W^C) + z F(W)$ and since $dF(w | w \in W) = dF(w) / F(W)$ we have that

$$\begin{aligned}
& \int_W E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}^\nu(\alpha; \theta), \theta, w) | s = \theta, \theta) dF(w | w \in W) \\
&= \int_{\bar{\alpha}(\alpha)}^{\bar{\alpha}(\alpha_P(\cdot, \theta))} u(z', \theta) \frac{dz'}{F(W)} + \int_W (v(\alpha_P(w, \theta), \theta, w) - v(a, \theta, w)) \frac{dF(w)}{F(W)} \\
&= \frac{1}{F(W)} (V(\alpha_P, \theta) - V(\alpha, \theta)) > 0. \tag{31}
\end{aligned}$$

Since $\alpha_P(w, \theta) > a$ for all $w \in W$ we have that $\bar{\alpha}(\alpha_P(\cdot, \theta)) > \bar{\alpha}(\alpha)$. Hence, by strict increasing differences we have that

$$\begin{aligned}
& E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}(\alpha_P(\cdot, \theta)), \theta, w) | s = \theta, \theta) \\
&> E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}^\nu(\alpha; \theta), \theta, w) | s = \theta, \theta).
\end{aligned}$$

Also, as $\nu \rightarrow 0$ expected payoff differences conditional on $s = \theta$ and θ converge to

the expected payoff differences conditional on only $s = \theta$. Accordingly,

$$\lim_{\nu \rightarrow 0} \int_W E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}(\alpha_P(\cdot, \theta)), \theta, w) | s = \theta) dF(w | w \in W) > 0. \quad (32)$$

The above argument applies to any $a > \alpha_P(w, \theta)$ by a symmetric application of increasing differences. Hence, inequality (32) is satisfied for any a and any subset W such that either $a < \alpha_P(w, \theta)$ or $a > \alpha_P(w, \theta)$, implying that

$$\lim_{\nu \rightarrow 0} E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}(\alpha_P(\cdot, \theta)), \theta, w) | s = \theta) > 0$$

for almost all w and all $a \neq \alpha_P(w, \theta)$. Since $\alpha_P(w, \cdot)$ is monotone it is continuous a.e. and the set of θ at which a positive mass of agents have a discontinuity in their strategies has measure zero. Hence, we have that $\lim_{\nu \rightarrow 0} \bar{\alpha}^\nu(\alpha_P; \theta) = \bar{\alpha}(\alpha_P(\cdot, \theta))$ for almost all $\theta \in (\inf \Theta, \sup \Theta)$, and the convergence is uniform. By the Lipschitz continuity of ΔU , this implies that

$$\lim_{\nu \rightarrow 0} E_\nu(\Delta U(\alpha_P(w, s), a, \bar{\alpha}^\nu(\alpha_P; \theta)), \theta, w) | s > 0$$

for almost all w and all $a \neq \alpha_P(w, \theta)$.

Finally, to prove the proposition under a general prior it suffices to show that (30) and (31) hold in the limit. This is because, in such a case, all the above arguments continue to apply when the prior is not uniform.

Note that the signal cutoff function is set to be $\kappa(w) = \theta$ for all W . Given such a cutoff function, we have that

$$\frac{\phi(\kappa(w))f(w)}{\int_W \phi(\kappa(w))f(w)dw} = \frac{f(w)}{\int_W f(w)dw} = f(w | w \in W).$$

Hence, applying Lemma 5 with $\kappa(w) = \theta$ for all W and taking the limit as $\nu \rightarrow 0$ we obtain (30) and (31). \square

B.5 Proofs of Results in Section 5

Proof of Lemma 2. If α^* is a NE with aggregate action \bar{a}^* then, for all a and w

$$\alpha^*(w)u(\bar{a}^*, \theta) + v(\alpha^*(w), \theta, w) \geq au(\bar{a}^*, \theta) + v(a, \theta, w).$$

Integrating these conditions across types we get that, for any strategy profile $\alpha \in \mathcal{A}$,

$$\bar{a}^*u(\bar{a}^*, \theta) + \int_w v(\alpha^*(w), \theta, w)dF(w) \geq \bar{\alpha}(\alpha)u(\bar{a}^*, \theta) + \int_w v(\alpha(w), \theta, w)dF(w). \quad (33)$$

But notice that, for any given aggregate action $\bar{a} \in [0, a_{max}]$ and any $\alpha \in \mathcal{A}_{\bar{a}}$, we have that $\int_w \alpha(w)u(\bar{a}, \theta)dF(w) = \bar{a}u(\bar{a}, \theta)$. Hence, by applying inequality (33) to the set of strategy profiles $\mathcal{A}_{\bar{a}^*}$ we obtain

$$\alpha^* \in \arg \max_{\alpha \in \mathcal{A}_{\bar{a}^*}} \int_w v(\alpha(w), \theta, w)dF(w). \quad (34)$$

Since a NE satisfies (33), given the definition of $B(\bar{a}, \theta)$, (34) implies that

$$\bar{a}^* \in \arg \max_{\bar{a} \in [0, a_{max}]} (\bar{a}u(\bar{a}^*, \theta) + B(\bar{a}, \theta)).$$

This proves the ‘‘if’’ part. To prove the ‘‘only if’’ part notice that if α^* satisfies (34) then it must yield the same average payoff as any NE profile associated with aggregate action \bar{a}^* . But then individual payoffs given \bar{a}^* must be maximized for all but a subset of types with measure zero, otherwise such NE would violate (33), and hence, the aggregate action of the two profiles must coincide. \square

Proof of Theorem 2. If $u(\cdot, \theta)$ is strictly increasing and $B(\cdot, \cdot)$ exhibits strictly increasing differences then, for any fixed \bar{a}^* the objective function in (14) also has strictly increasing differences in \bar{a} and θ . In addition, it is straightforward to check that the objective function is continuous in \bar{a} and A is compact.²² Hence, by Topkis monotonicity theorem the correspondence

$$m(\bar{a}^*, \theta) := \arg \max_{\bar{a} \in [0, a_{max}]} (\bar{a}u(\bar{a}^*, \theta) + B(\bar{a}, \theta))$$

is increasing in θ in the strong set order. In addition, $m(\bar{a}^*, \theta)$ is non-empty and upper hemicontinuous by Berge’s maximum theorem.

Finally, notice that $\min m(0, \theta) \geq 0$ and $\max m(a_{max}, \theta) \leq a_{max}$ for all θ . That is, the correspondence $m(a_{max})$ ‘crosses’ the diagonal from above at the smallest and largest crossing values of \bar{a}^* . But since the NE aggregate actions satisfy $\bar{a}^* \in m(\bar{a}^*, \theta)$, i.e., they are represented by the values at which $m(\cdot, \theta)$ crosses the diagonal, the smallest and largest crossing values must go up as $m(\cdot, \theta)$ increases with θ . \square

Proof of Theorem 3. Given any $\alpha \in \text{int}(\mathcal{A})$, consider an infinitesimal increase da of all individual actions $\alpha(w)$. Since v is Lipschitz continuous it is differentiable almost everywhere, the change in average idiosyncratic payoffs for almost all $\alpha \in \text{int}(\mathcal{A})$ is given by

$$\int_w \frac{\partial v(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} dF(w).$$

²²By Proposition 2 a NE always exists implying that the maximization problem associated with $B(\bar{a}, \theta)$ has a solution. In addition, the continuity of B follows from v being continuous and bounded and the fact that, for any $\alpha \in \mathcal{A}_{\bar{a}}$, we can find a profile $\alpha' \in \mathcal{A}_{\bar{a}+\varepsilon}$ such that $|\alpha'(w) - \alpha(w)| = \varepsilon$ and vice versa for all w .

But notice that such an increase in individual actions implies that the aggregate action also increases by da . Hence, we must have that

$$\frac{\partial B(\bar{a}, \theta)}{\partial \bar{a}} \geq \max_{\alpha \in \text{int}(\mathcal{A}_{\bar{a}})} \int_{\underline{w}}^{\bar{w}} \frac{\partial v(a, \theta, w)}{\partial a} \Big|_{a=\alpha(w)} dF(w).$$

Accordingly, we must have that

$$\frac{\partial^2 B(\bar{a}, \theta)}{\partial \bar{a} \partial \theta} \geq \max_{\alpha \in \text{int}(\mathcal{A}_{\bar{a}})} \int_{\underline{w}}^{\bar{w}} \frac{\partial^2 v(a, \theta, w)}{\partial a \partial \theta} \Big|_{a=\alpha(w)} dF(w) > 0.$$

That is, $B(\bar{a}, \theta)$ exhibits strictly increasing differences in $(0, a_{max}) \times \Theta'$ and, by continuity, in $[0, a_{max}] \times \Theta'$ so [Theorem 2](#) applies. \square

Proof of [Theorem 4](#). The proposition immediately follows by noting that (i) the first term in the objective function of [\(14\)](#) does not depend on F and, (ii) we can index F and \hat{F} using parameter $\zeta \in \mathbb{R}$ so that the condition in the proposition is equivalent to B exhibiting increasing differences in \bar{a} and ζ . Together, (i) and (ii) imply that u and B satisfy the conditions in [Theorem 2](#) w.r.t. \bar{a} and ζ (keeping θ unchanged). \square

Proof of [Theorem 5](#). The proof logic is similar to the one used in [Theorem 3](#) and is therefore omitted. \square

References

- Abadi, Joseph and Markus Brunnermeier**, “Blockchain economics,” 2018. Working Paper.
- Acemoglu, Daron and Martin Kaae Jensen**, “Robust Comparative Statics in Large Static Games,” in “49th IEEE Conference on Decision and Control (CDC)” IEEE 2010, pp. 3133–3139.
- **and –**, “Robust Comparative Statics in Large Dynamic Economies,” *Journal of Political Economy*, 2015, *123* (3), 587–640.
- Bassetto, Marco and Christopher Phelan**, “Tax Riots,” *Review of Economic Studies*, 2008, *75* (3), 649–669.
- Basteck, Christian, Tijmen R Daniëls, and Frank Heinemann**, “Characterising Equilibrium Selection in Global Games with Strategic Complementarities,” *Journal of Economic Theory*, 2013, *148* (6), 2620–2637.
- Bond, Philip and Kathleen Hagerty**, “Preventing Crime Waves,” *American Economic Journal: Microeconomics*, 2010, *2* (3), 138–159.

- Camacho, Carmen, Takashi Kamihigashi, and Çağrı Sağlam**, “Robust Comparative Statics for Non-monotone Shocks in Large Aggregative Games,” *Journal of Economic Theory*, 2018, *174*, 288–299.
- Carlsson, Hans and Eric van Damme**, “Global Games and Equilibrium Selection,” *Econometrica*, September 1993, *61* (5), 989–1018.
- Cheung, Man-Wah and Ratul Lahkar**, “Nonatomic Potential Games: the Continuous Strategy Case,” *Games and Economic Behavior*, 2018, *108*, 341–362.
- Cornes, Richard and Roger Hartley**, “Asymmetric Contests with General Technologies,” *Economic Theory*, 2005, *26* (4), 923–946.
- Dasgupta, Partha S and Geoffrey M Heal**, *Economic Theory and Exhaustible Resources*, Cambridge University Press, 1979.
- Diamond, Douglas W and Philip H Dybvig**, “Bank Runs, Deposit Insurance, and Liquidity,” *Journal of Political Economy*, 1983, *91* (3), 401–419.
- Diamond, Peter A**, “Aggregate Demand Management in Search Equilibrium,” *Journal of Political Economy*, 1982, *90* (5), 881–894.
- Drozd, Lukasz A. and Ricardo Serrano-Padial**, “Financial Contracting with Enforcement Externalities,” *Journal of Economic Theory*, 2018, *178*, 153–189.
- Dubey, Pradeep, Ori Haimanko, and Andriy Zapechelnyuk**, “Strategic Complements and Substitutes, and Potential Games,” *Games and Economic Behavior*, 2006, *54* (1), 77–94.
- Dybvig, Philip H and Chester S Spatt**, “Adoption Externalities as Public Goods,” *Journal of Public Economics*, 1983, *20* (2), 231–247.
- Farrell, Joseph and Garth Saloner**, “Standardization, Compatibility, and Innovation,” *The RAND Journal of Economics*, 1985, pp. 70–83.
- Frankel, David M., Stephen Morris, and Ady Pauzner**, “Equilibrium Selection in Global Games with Strategic Complementarities,” *Journal of Economic Theory*, 2003, *108* (1), 1–44.
- Goldstein, Itay and Ady Pauzner**, “Demand–deposit Contracts and the Probability of Bank Runs,” *Journal of Finance*, 2005, *60* (3), 1293–1327.
- Guimaraes, Bernardo and Stephen Morris**, “Risk and Wealth in a Model of Self-fulfilling Currency Attacks,” *Journal of Monetary Economics*, 2007, *54* (8), 2205–2230.

- Hofbauer, Josef and William H Sandholm**, “Evolution in Games with Randomly Disturbed Payoffs,” *Journal of Economic Theory*, 2007, 132 (1), 47–69.
- Jensen, Martin Kaae**, “Aggregative Games,” in L.C. Corchón and M. A. Marini, eds., *Handbook of Game Theory and Industrial Organization*, Edward Elgar, 2018.
- Kajii, Atsushi and Stephen Morris**, “The Robustness of Equilibria to Incomplete Information,” *Econometrica*, 1997, pp. 1283–1309.
- Katz, Michael L and Carl Shapiro**, “Technology Adoption in the Presence of Network Externalities,” *Journal of Political Economy*, 1986, 94 (4), 822–841.
- Lahkar, Ratul**, “Large Population Aggregative Potential Games,” *Dynamic Games and Applications*, 2017, 7 (3), 443–467.
- Martimort, David and Lars Stole**, “Representing Equilibrium Aggregates in Aggregate Games with Applications to Common Agency,” *Games and Economic Behavior*, 2012, 76 (2), 753–772.
- Milgrom, Paul and Chris Shannon**, “Monotone Comparative Statics,” *Econometrica*, 1994, 62 (1), 157–180.
- **and John Roberts**, “Rationalizability, Learning, and Equilibrium in Games with Strategic Complementarities,” *Econometrica*, 1990, 58 (6), 1255–1277.
- Monderer, Dov and Lloyd S Shapley**, “Potential games,” *Games and economic behavior*, 1996, 14 (1), 124–143.
- Morris, Stephen and Hyun Song Shin**, “Unique Equilibrium in a Model of Self-Fulfilling Currency Attacks,” *American Economic Review*, 1998, 88 (3), pp. 587–597.
- **and** – , “Global Games: Theory and Applications,” in “Proceedings of the Eighth World Congress of the Econometric Society,” Cambridge University Press, 2003.
- **and Takashi Ui**, “Generalized Potentials and Robust Sets of Equilibria,” *Journal of Economic Theory*, 2005, 124 (1), 45–78.
- Moulin, Herve**, “Joint Ownership of a Convex Technology: Comparison of Three Solutions,” *The Review of Economic Studies*, 1990, 57 (3), 439–452.
- Obstfeld, Maurice**, “Rational and Self-fulfilling Balance-of-Payments Crises,” *American Economic Review*, 1986, 76 (1), 72–81.
- Oyama, Daisuke and Satoru Takahashi**, “Generalized Belief Operator and Robustness in Binary-Action Supermodular Games,” *Econometrica*, 2020, 88 (2), 693–726.

- Sakovics, Jozsef and Jakub Steiner**, “Who Matters in Coordination Problems?,” *American Economic Review*, 2012, 102 (7), 3439–3461.
- Sandholm, William H**, “Large Population Potential Games,” *Journal of Economic Theory*, 2009, 144 (4), 1710.
- Serrano-Padial, Ricardo**, “Coordination in Global Games with Heterogeneous Agents,” 2018. Working Paper.
- Topkis, Donald M**, “Equilibrium Points in Nonzero-sum N-person Submodular Games,” *Siam Journal on control and optimization*, 1979, 17 (6), 773–787.
- Tullock, Gordon**, “Efficient Rent Seeking,” in G. Tullock JM Buchanan, RD Tol-
lison, ed., *Toward a Theory of the Rent Seeking Society*, Texas A & M University
Press, 1980, chapter 4.
- Ui, Takashi**, “Robust Equilibria of Potential Games,” *Econometrica*, 2001, 69 (5),
1373–1380.
- Vives, Xavier**, “Nash Equilibrium with Strategic Complementarities,” *Journal of
Mathematical Economics*, 1990, 19 (3), 305–321.
- Zandt, Timothy Van and Xavier Vives**, “Monotone Equilibria in Bayesian
Games of Strategic Complementarities,” *Journal of Economic Theory*, 2007, 134
(1), 339–360.
- Zusai, Dai**, “Evolutionary Dynamics in Heterogeneous Populations: A General
Framework without Aggregability,” 2018. Working Paper.